

Indeks funkcionalnih skupina fitoplanktona kao pokazatelj promjena ekološkog stanja jezera u poplavnom području Dunava

Kajan, Katarina

Master's thesis / Diplomski rad

2017

Degree Grantor / Ustanova koja je dodijelila akademski / stručni stupanj: **Josip Juraj Strossmayer University of Osijek, Department of biology / Sveučilište Josipa Jurja Strossmayera u Osijeku, Odjel za biologiju**

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:181:729860>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-03-29**



Repository / Repozitorij:

[Repository of Department of biology, Josip Juraj Strossmayer University of Osijek](#)



**JOSIP JURAJ STROSSMAYER UNIVERSITY OF OSIJEK
DEPARTMENT OF BIOLOGY**

Graduate university study programme in Biology

Katarina Kajan

**PERFORMANCE OF THE PHYTOPLANKTON
ASSEMBLAGE INDEX IN EVALUATION OF
ENVIRONMENTAL CHANGES OF A DANUBIAN
FLOODPLAIN LAKE**

Master's Thesis

OSIJEK, 2017.

TEMELJNA DOKUMENTACIJSKA KARTICA

Sveučilište Josipa Jurja Strossmayera u Osijeku
Odjel za biologiju
Diplomski sveučilišni studij Biologija; smjer: znanstveni

Diplomski rad

Znanstveno područje: Prirodne znanosti
Znanstveno polje: Biologija

INDEKS FUNKCIONALNIH SKUPINA FITOPLANKTONA KAO POKAZATELJ PROMJENA EKOLOŠKOG STANJA JEZERA U POPLAVNOM PODRUČJU DUNAVA

Katarina Kajan

Rad je izrađen: Zavod za ekologiju voda, Odjel za biologiju, Sveučilište Josipa Jurja Strossmayera u Osijeku

Mentor: dr. sc. Melita Mihaljević, izvanredni profesor

Komentor: dr. sc. Judit Padisák, redoviti profesor, Odjel za limnologiju, Sveučilište Pannonia, Veszprém, Mađarska

Indeks funkcionalnih skupina fitoplanktona (Q indeks) koristi se kao pokazatelj promjena ekološkog stanja slatkovodnih sustava te se istražuju mogućnosti njegove primjene u ocjeni stanja različitih tipova jezera. Cilj ovog rada je na temelju dinamike fitoplanktonskih funkcionalnih skupina i primjenom Q indeksa ocijeniti ekološko stanje Sakadaškog jezera, plitkog jezera u sustavu poplavnog područja Dunava. Istraživanje je provedeno u razdoblju srpanj 2011-listopad 2012, u hidrološkim uvjetima uobičajene dinamike poplava. Rezultati su uspoređeni s povijesnim podatcima dinamike fitoplanktona u razdoblju srpanj 1972-rujan 1973, kada je jezero bilo pod jakim antropogenim utjecajem. Dinamika i zastupljenost pojedinih funkcionalnih skupina fitoplanktona značajno su se razlikovali između dva uspoređena razdoblja. Česta pojava masovnog razvoja jedne vrste ili funkcionalne skupine karakteristične za zagađene vode, bile su značajke dinamike fitoplanktona u uvjetima velikog zagađenja okoliša na što ukazuju niske vrijednosti Q indeksa i ocjena *loše* do *umjereno loše* stanje. U uvjetima bez antropogenih pritisaka i očuvanog hidrološkog stanja, u fitoplanktonskoj su zajednici bile zastupljene funkcionalne skupine karakteristične za prirodna eutrofna jezera, a vrijednosti Q indeksa pokazuju da je jezero u statusu između *umjerenog* i *dobrog* stanja. Pomoću Q indeksa mogu se vrlo precizno odrediti promjene ekološkog statusa jezera.

Broj stranica: 52

Broj slika: 14

Broj tablica: 4

Broj literaturnih navoda: 95

Broj priloga: 1

Jezik izvornika: engleski

Ključne riječi: jezero u poplavnom području, funkcionalne skupine fitoplanktona, Q indeks, ekološki status, antropogeni utjecaji

Datum obrane: 31.10.2017.

Stručno povjerenstvo za obranu:

1. Dr. sc. *Dubravka Špoljarić Maronić*, docent, predsjednik povjerenstva
2. Dr. sc. *Melita Mihaljević*, izvanredni profesor, mentor, član
3. Dr. sc. *Judit Padisák*, redoviti profesor, komentor, član
4. Dr. sc. *Filip Stević*, docent, zamjena člana

Rad je pohranjen u: knjižnici Odjela za biologiju Sveučilišta Josipa Jurja Strossmayera u Osijeku i u Nacionalnoj i sveučilišnoj knjižnici u Zagrebu, u elektroničkom obliku te je objavljen na web stranici Odjela za biologiju

BASIC DOCUMENTATION CARD

Josip Juraj Strossmayer University of Osijek
Department of Biology
Graduate university study programme in Biology
Scientific Area: Natural science
Scientific Field: Biology

Master's Thesis

PERFORMANCE OF THE PHYTOPLANKTON ASSEMBLAGE INDEX IN EVALUATION OF ENVIRONMENTAL CHANGES OF A DANUBIAN FLOODPLAIN LAKE

Katarina Kajan

Thesis performed at Subdepartment of Water Ecology, Department of Biology, Josip Juraj Strossmayer University of Osijek

Supervisor: Melita Mihaljević, PhD, Associate Professor

Co-supervisor: Judit Padisák, PhD, Full Professor, Department of Limnology, University of Pannonia, Veszprém, Hungary

As a monitoring tool for ecological status assessment of freshwater bodies, phytoplankton assemblage index (Q index) was so far successfully applied in the evaluation of different types of lakes. Currently, its wider application possibilities are being investigated. The aim of this study was to evaluate ecological status of the floodplain lake, Lake Sakadaš, part of the complex river-floodplain system in the Middle Danube (Croatia), by assessment of phytoplankton functional groups and applying the Q index. Phytoplankton research was conducted in the period July 2011-October 2012, in conditions of more or less usual flooding pattern. Results were compared with the historical data of phytoplankton dynamics within the period July 1972-September 1973, when the lake was under the strong anthropogenic pressure, which resulted in rapid eutrophication and deterioration of its ecological status. The data set of phytoplankton functional groups and the main group arrangements in the multivariate nMDS analysis showed variation related to the past and recent conditions. The frequent appearance of heavy bloom of only one species or assemblages characteristic for polluted waters was the most distinctive feature of phytoplankton dynamics in the past. In such an impacted environment, values of the Q index were low and indicated *bad* to *tolerable* ecological status. Recent data suggest that water quality improvement and near-natural hydrological condition support algal assemblages characteristic for naturally eutrophic lakes and the values of the Q index varied between *medium* to *good* ecological status. Altogether, results of this study showed that Q index can very precisely reflect anthropogenic pressure what was recognized as its major advantage in comparison with other indices used for ecological status assessment.

Number of pages: 52

Number of figures: 14

Number of tables: 4

Number of references: 95

Number of appendices: 1

Original in English

Keywords: floodplain lake, phytoplankton functional groups, Q index, ecological status, anthropogenic pressure

Date of the thesis defense: 31st October 2017

Reviewers:

1. *Dubravka Špoljarić Maronić*, PhD, Assistant Professor
2. *Melita Mihaljević*, PhD, Associate Professor, supervisor
3. *Judit Padisák*, PhD, Full Professor, co-supervisor
4. *Filip Stević*, PhD, Assistant Professor

Thesis deposited in: Library of Department of Biology, University of J. J. Strossmayer Osijek and in a National and University Library in Zagreb in electronic form. It is also disposable on the website of Department of Biology, University of J. J. Strossmayer Osijek.

Acknowledgements

I express my utmost gratitude to my supervisor Prof. Dr. Melita Mihaljević for her invaluable guidance, constructive comments and helpful suggestions which substantially improved this thesis.

I would like to express my sincerest appreciativeness to my co-supervisor Prof. Dr. Judit Padisák from Department of Limnology at the University of Pannonia in Veszprém, on the opportunity to do my student training in the department. Her expert guidance and many helpful suggestions helped me tremendously in the process of writing this thesis.

I am truly grateful to Dr. Géza Szelmeczy for his support and guidance during my Erasmus student training in Department of Limnology.

I would also like to acknowledge the staff of Laboratory of Ecology of Algae, particulary to Dr. Dubravka Špoljarić Maronić and Dr. Tanja Žuna Pfeiffer on their support and availability whenever I needed help during my work in the laboratory.

A special thank to the family of Prof. Dr. Dragica Gucunski for encouraging us in proccesing of historical data.

I thank to my copy editor Martina Leko for careful reading of the thesis and language corrections.

Finally, I express my profound gratitude to my family for providing me with their unfailing support and continuous encouragement throughout my years of study.

TABLE OF CONTENTS

1. INTRODUCTION	1
1.1. Study aim	4
2. MATERIALS AND METHODS	5
2.1. Study area	5
2.2. Sampling and analysis of physical and chemical parameters	8
2.3. Phytoplankton analyses	9
2.4. Data analysis	10
2.5. Statistical analyses	10
3. RESULTS	11
3.1. Physical and chemical characterization of the environment	11
3.2. Taxonomical and functional diversity of phytoplankton	18
3.3. Phytoplankton assemblage index	26
4. DISCUSSION	29
5. CONCLUSIONS	34
6. REFERENCES	35
7. APPENDIX	44

1. INTRODUCTION

The ecological status estimation of surface waters according to the Water Framework Directive (Directive 2000/60/EC; WFD, 2000) is based on the assessment of biological elements supported by hydromorphological, chemical and physicochemical elements. The phytoplankton as an extremely diverse, photoautotrophic group of organisms responsible for the majority of primary oxygen and organic carbon production in aquatic ecosystems (Reynolds, 2006) is one of the five biological quality elements proposed for ecological status assessment of surface waters. Phytoplankton species are good indicators of water quality because they reflect the changes very rapidly both in their quantitative and qualitative composition. Dynamics of phytoplankton are mainly determined by environmental changes occurring at different temporal scales (short-term and annual variations) and are regulated by both internal and external factors (Reynolds, 1993; Reynolds, 2006).

Traditional phytoplankton monitoring based on the taxonomic level of community structure, including phytoplankton biomass and chlorophyll *a* concentration, appeared to be only partially useful for determining water quality. The main disadvantage of this approach is that numerous species of phytoplankton are included in the broader taxonomic groups despite their very different ecological properties (Reynolds et al., 2002).

The monitoring of phytoplankton was supplemented with phytoplankton trait-based approaches which grouped species with similar morphological and functional properties. Several phytoplankton classification concepts were developed: the functional group (FG) classification (Reynolds et al., 2002; Padisák et al., 2009), the morpho-functional group (MFG) classification (Salmaso and Padisák, 2007) and morphology-based group (MBFG) classification (Kruk et al., 2010; Kruk and Segura, 2012).

According to physiological, morphological and ecological attributes of phytoplankton species, Reynolds et al. (2002) proposed an approach assigning phytoplankton species into functional groups (coda). Groups of species are defined with specific habitat, tolerances and sensitivities based on several different combinations of physical, chemical and biological properties of the water body environment (such as depth of mixing layer, light, temperature, P, N, Si, CO₂, grazing pressure). The approach was further developed by Padisák et al. (2009) stipulating a detailed description of the typical misplacements and by modifying some of the original habitat templates and species allocations. Functional classification, altogether with more than 40 groups (coda) described, simplifies the comparison of seasonal changes in various water body

types and evaluates the responses to environmental changes (Kruk et al., 2002; Naselli-Flores et al., 2003). The classification has been applied worldwide from temperate to tropical regions in different ecosystems, particularly in shallow lakes (Huszar et al., 2003; Mieleitner et al., 2008; Rangel et al., 2009; Barone et al., 2010; Borics et al., 2012; Izaguirre et al., 2012; Crossetti et al., 2013; Hu et al., 2013). This classification was also shown as successful approach in monitoring the changes of phytoplankton in reservoirs (Fonseca and Bicudo, 2008; Borges et al., 2008; Becker et al., 2009a; Xiao et al., 2011), estuaries (Costa et al., 2009) and rivers (Borics et al., 2007; Abonyi et al., 2012; Stanković et al., 2012).

Following the functional group classification, Padisák et al. (2006) developed the phytoplankton assemblage index (Q index) initially proposed to assess the ecological status of different lake types in line with the WFD (2000). The assemblage index provides 5 degrees of water quality based on the relative share of FG multiplied by factor number (F) defined for each FG. The most crucial step of the assessment is a determination of factor F values because they reflect values of FG in the reference condition of the studied water body (pristine status). The Q index was initially developed to assess the ecological status of different Hungarian lake types (Padisák et al., 2006). According to that time existing typology, there are 8 lake types in Hungary to which belong large lakes, alkaline lakes and oxbows outside flood control dams. Values of factor F were established for each functional group within all lake types.

Hajnal and Padisák (2008) used the phytoplankton biomass and assemblage structure expressed by the Q index to reconstruct the history of water quality in Lake Balaton and to quantify the changes in water quality during the eutrophication and restoration phases. This research represents the first attempt at historical reconstruction of the ecological status and compares it with changes in trophic state and current water quality. The results showed that this method could be successfully used to reconstruct water quality from historical archives and it is helpful in defining reference condition of the water body.

Up to now, the Q index was successfully applied to evaluate ecological status of various types of lakes (Demir et al., 2014; Teneva et al., 2014; Ochocka and Pasztaleniec, 2016; Ongun Sevindik et al., 2017) and other types of water bodies such as reservoirs (Crossetti and Bicudo, 2008; Becker et al., 2009b; Becker et al., 2010; Wang et al., 2011; Çelekli and Öztürk, 2014; Molina-Navarro et al., 2014; Vieira et al., 2015; Silva and Costa, 2015; Santana et al., 2017) regardless of geographic regions. It should be emphasized that the Q index was successfully applied for assessment of the water bodies from existed lake types in Bulgaria according to the accepted typology (Belikinova et al., 2014).

Floodplain lakes, also known as oxbow lakes, fluvial lakes, river lakes, limnocren, crescent lakes (Dawidek and Ferencz, 2012 and cites therein) remain to be the least known group of water bodies due to their complexity and diversity. A floodplain lake includes every inland water body, whose basin originates from fluvial processes, and its limnological functioning derives from irregular, but periodical limnophases and potamophases (Dawidek and Ferencz, 2012). Characterization of a floodplain lake is based on hydrogeological, hydrological and hydrochemical features. Due to the high anthropogenic impact, which includes the alteration of natural flow pulses, channelization, dredging, artificial levee construction, groundwater abstraction and enrichment with nutrients (wastewaters, nutrient inflow from agricultural areas), floodplains are nowadays one of the most threatened and sensitive ecosystems (Amoros and Bornette, 2002). The majority of large active floodplains have disappeared over the last two centuries, including up to 80% of the historical floodplain area along the Danube and its larger tributaries (Schwarz, 2010).

Floodplain along the Middle Danube section, known as Kopački Rit Nature Park, is one of the largest preserved natural floodplains of the Danube River. A diversity of aquatic and wet biotopes composed of ephemeral and perennial water bodies, mostly channels, oxbows, marshes and shallow lakes, continuously changing the area covered by water depending on the inflow of riverine waters. One of the significant water bodies in this floodplain is Lake Sakadaš, the deepest water depression located in the western part of the floodplain. During the 1970s, when the first limnological investigations were carried out in the Lake Sakadaš (primarily focused on the water quality and phytoplankton), high variability in physical and chemical parameters was detected by water quality assessment. High variabilities in parameters indicated the high anthropogenic impact on the lake due to the direct income of wastewaters from the surrounding agricultural areas, which resulted in rapid eutrophication and deterioration of the ecosystem (Gucunski, 1975). Efforts for protection of the whole floodplain habitats resulted in strong protection of the area with the status of Nature Park. Moreover, measures for the revitalization of Lake Sakadaš applied in 1984 included complete isolations from the incoming of wastewaters and sediment removal.

During the past few decades, the lake was in eutrophic/hypertrophic state according to the OECD system (OECD, 1982) with yearly average of TP concentration higher than $100 \mu\text{g L}^{-1}$, water transparency less than 1.5 m, and maximum chlorophyll *a* concentration higher than $50 \mu\text{g L}^{-1}$ (Mihaljević et al., 1999; Vidaković and Bogut, 2007; Horvatić et al., 2003; Stević et al., 2005). The results of permanent monitoring of phytoplankton in the lake were carried out from

2003 till nowadays, showed that depending on the time scale occurrence, flood pulses can be a stimulating or a disturbance factor for phytoplankton development in Lake Sakadaš (Mihaljević et al., 2009). However, the intensity and duration of flooding can be qualified as the primary cause for the changes of phytoplankton assemblages in this floodplain lake (Stević et al., 2013).

1.1. Study aim

The aim of this study was to evaluate ecological status of the floodplain lake, Lake Sakadaš applying phytoplankton assemblage index. Due to the fact that there are no historical data for evaluation of pristine condition in this floodplain lake, current research of phytoplankton assemblages dynamic will be compared with the historical data of phytoplankton dynamics when the lake was under strong anthropogenic pressure. The hypothesis is that Q index will indicate changes between impacted and semi-natural conditions of the floodplain lake.

2. MATERIALS AND METHODS

2.1. Study area

Kopački Rit Nature Park is one of the largest conserved natural riverine floodplains located in North-East Croatia between the Drava and the Danube River (Figure 1.). It is internationally recognized as the Ramsar Area (No: 3HR002) protected by the Ramsar Convention and included in the Important Bird Area (IBA) list, covering approximately 180 km². Since the Pleistocene and Holocene epochs, when the Kopački Rit was formed, its ecological equilibrium and existence depends on the flooding regime of Danube and Drava River (Tadić et al., 2014). The floodplain complex provides a diversity of biotopes composed of periodic and permanent water bodies, such as marshes/pools, ponds, swamps on organic soils, shallow lakes, riverside arms and natural channels. The hydrological dynamics of the floodplain is under the impact of horizontal (inflows and outflows) and vertical (precipitation, evaporation and transpiration) components of water balance (Tadić et al., 2014).

The Drava River has a minor influence due to the high embankments constructed in the middle of the 20th century between the river and the floodplain (Bonacci et al., 2002). The intensity of the flooding primarily depends on the hydrological regime of the Danube, and it may occur in any season of the year. In the middle section, the Danube (1,410- 1,383 r. km) is a typical lowland river with the mean annual water level of 2.63 m and mean annual discharge of 2,085 m³s⁻¹ (from a database of daily recordings in the period 1987-2008 at the gauge station at river 1,401.4 km). Generally, during the first half of the year (mid-spring), the flow of the Danube is the highest, while in the second half of the year (June-October) the flow is characterized by a decrease (Buijse et al., 2002). During low river water discharge, the permanent water biotopes in the floodplain are isolated from one another.

The floodplain area can be divided into two subsystems due to the hydrological connectivity with the main river channel. Subsystem A is impounded by the river through the backwater system (side arm), while subsystem B through a network of perennial channels. According to Schwarz (2005), minor floods (3-3.5 m) inundate only 18% of the area in subsystem B, while extremely high floods (more than 5 m) inundate almost the whole floodplain area (more than 90% of subsystem B).




Figure 1. Location of the sampling site Lake Sakadaš in the study area Nature Park Kopački Rit. The dotted line marks the border of Nature Park Kopački Rit and dashed line marks the embankment of rivers Danube and Drava (arrows indicate the flow direction). Modified from Mihaljević et al. (2013).

The study was carried out in the western part of the floodplain in the Lake Sakadaš (Figure 1., Figure 2.). The average depth of the lake is about 5 m and a maximum depth of ca. 12 m, with a surface water area of about 0.15 km². It is in direct hydrological connection with the main river channel through two channels (Čonakut and Hulovo channels) with total length ca. 10 km, and it is close to the embankment which delineates the inundation area. Flooding of the lake begins when the Danube water level at the gauge station river 1,401.4 km rises above 3 m (Mihaljević et al., 1999). The mixing regime is closely connected to flood events, and thermal stratification is usually expressed during the summer (June-September), while during the winter months (December-February) the entire lake is ice-covered.

In shallow parts of the lake, submerged macrophytes are well developed (*Ceratophyllum demersum* L., *Myriophyllum spicatum* L. and *Potamogeton gramineus* L.). The common reed (*Phragmites communis* Trin.) occurs around the lakeshore adjoined by floodplain forests of white willow (*Salix alba* L.) and black poplar (*Populus nigra* L.).



Figure 2. Sampling site Lake Sakadaš. (Pictures from Subdepartment of Water Ecology archives)

2.2. Sampling and analysis of physical and chemical parameters

Sampling was conducted at monthly intervals during the period from July 2011 to October 2012 (except in February 2012) in the central part of Lake Sakadaš. The main studied physical parameters were water temperature, water depth, transparency, conductivity, pH and dissolved oxygen. Water temperature, pH, conductivity and dissolved oxygen were measured *in situ* at subsurface (*ca.* 0.2 m) using portable instrument Multi 340i WTW (Wissenschaftlich-Technische Werkstätten, Weilheim, Germany). Water depth of the lake was measured using a weighted rope, and the transparency was estimated with a Secchi disc. Samples for the chemical analyses of ammonium (NH_4^+), nitrates (NO_3^-), nitrites (NO_2^-), organic nitrogen (orgN), total nitrogen (TN) and total phosphorus (TP) were taken at a depth of *ca.* 0.2 m below the water surface. Chemical variables were analyzed in the laboratory according to standard methods (APHA, 1992). In order to determine the concentration ($\mu\text{ l}^{-1}$) of chlorophyll *a* (Chl *a*), *-b* (Chl *b*) and *-c* (Chl *c*), sample of 1-L was carried on ice to the laboratory and filtered through Whatman GF/C glass fibre filters (Whatman International Ltd., Maidstone, England). Filtered samples were subsequently extracted with acetone. Absorbance was measured with a Hach DR 2010 spectrophotometer (Hach Company, USA) at four different wavelengths (630, 645, 663 and 750 nm). Chl *a*, *-b* and *-c* concentrations were calculated according to UNESCO (1966) and Strickland and Parsons (1972).

Similar methods were applied during the period from July 1972 to September 1973 (except in August 1973), with data available in the doctoral thesis of Professor Dragica Gucunski (Gucunski, 1975). From the doctoral thesis, the following values of physical variables were available: water temperature, water depth, transparency, pH and dissolved oxygen. Values of chemical variables could not be used, wherein only their presence or absence is indicated.

Daily records of Danube water level were obtained for the gauge station at river 1,401.4 km.

2.3. Phytoplankton analyses

Depth-integrated samples intended to assess the quantitative composition of phytoplankton were collected from the entire water column and fixed *in situ* with Lugol's acidified solution. Additional samples for qualitative taxonomic analysis were obtained with a 22.5 µm pore net and fixed with 4% formalin. Phytoplankton species were identified with light microscopic observations (Carl Zeiss Jena, Jena, Germany) using the standard literature for species determination (Hustedt, 1976; Hindák et al., 1978; Meffert et al., 1981; Anagnostidis and Komárek, 1985, 1988; Komárek and Anagnostidis, 1989). The taxonomic classification was updated with recent findings, and the nomenclature was updated according to Algaebase website (Guiry and Guiry, 2014). Quantitative assessment of phytoplankton was done according to Utermöhl (1958) using the inverted microscope (Axiovert 25, Carl Zeiss, Inc., Göttingen, Germany) at multiple magnifications (100×, 400×) counting 400 individuals. The counting unit was the individual (unicell, coenobium, filament or colony). Each species abundance was presented as the number of individuals per liter (ind. L⁻¹). Biovolumes were calculated following Rott (1981) where individuals were measured and their volumes calculated by relating cell shape with the corresponding geometric body. Volume calculations of colonial organisms with mucilage included entire colonies together with mucilage. Biomass was estimated multiplying each phytoplankton species abundance with mean biovolume of species (Javornický and Komárková, 1973; Sournia, 1978) and expressed as milligrams per liter (mg l⁻¹) fresh mass.

Historical data of phytoplankton quantitative analysis from July 1972-September 1973 was used from the doctoral thesis of Professor Dragica Gucunski (Gucunski, 1975) with her family's permission. Data of phytoplankton abundance (cell L⁻¹) was converted to biomass (mg L⁻¹), calculated according to cell biovolume of phytoplankton species database published by Gucunski and Popović (1984). Historical phytoplankton data was updated with currently used taxonomic classification and nomenclature.

2.4. Data analysis

Each taxon, determined in the study period from July 2011 to October 2012 and in historical data from July 1972 to September 1973, was classified into the functional group proposed in the classification of Reynolds et al. (2002) revised by Padisák et al. (2009). The assemblage index (Q index) followed by Padisák et al. (2006) was applied to assess the ecological status of floodplain Lake Sakaš during the periods.

The assemblage index was calculated according to the following formula:

$$Q = \sum_{i=1}^n p_i F$$

p_i – relative share of functional groups in total biomass
 $p_i = n_i/N$ n_i : biomass of the i -th functional group
 N : total biomass
 F – factor number established for the i -th functional group

The relative share of FGs in total biomass was calculated, and factor numbers (F) were established for each FG registered in chosen periods. Factor number was determined in the range from 0 to 5. Determination of factor numbers was based on the previous knowledge and experience of experts, together with the literature data of phytoplankton composition in Lake Sakadaš from 1972 up to the present day. Given factor numbers were compared with F defined for similar water body types (Padisák et al., 2006; Crossetti et Bicudo, 2008; Becker et al., 2010; Belkinova et al., 2014). According to the WFD requirements, values of Q index between 0 and 1 were classified as bad, between 1 and 2 as tolerable, between 2 and 3 as medium, between 3 and 4 as good, and between 4 and 5 as excellent (Padisák et al., 2006).

2.1. Statistical analyses

Pearson's correlation coefficient was used to analyze the correlation between the physical and chemical parameters of the periods using Statistica 8.0 software (StatSoft, Inc., USA).

Non-metric multidimensional scaling (nMDS) was used to display similarity between the biomass of phytoplankton FGs of chosen periods using the statistical program PRIMER version 5.0 (Clarke and Warwick, 2001). Biomass of phytoplankton FGs was the square root transformed and applied on Bray-Curtis similarity coefficient. Increasing distance between samples in nMDS plot reflects higher dissimilarities between the biomass of FGs in the periods.

3. RESULTS

3.1. Physical and chemical characterization of the environment

Flooding dynamics of the river-floodplain system is determined by the fluctuation of the Danube water level. Daily courses of the Danube water level at river 1,401.4 km are shown in Figure 3. During the periods of July 1972-October 1973 and July 2011-October 2012, the floodplain was characterized by the periodical occurrence of limnophase (dry periods) and potamophase (flood periods). Flooding patterns were defined by the timing (season), duration of flooding and the total surface of the flooded area (Table 1.).

Danube water level daily courses of chosen periods showed a statistically significant positive correlation ($r = 0.27$, $p < 0.05$), despite the higher maximal water level (5.84 m) and extremely high flooding (27 days in total) with more than 75% of the flooded area recorded in 1972-1973. Most extended flood with 81 days in continuum was present during the spring-summer period of 1973. The absence of flooding with 155 days in total characterized the longest dry phase of studied period 2011-2012. After the dry phase, short-time flood pulses were frequent during the late winter and spring with two major flood pulses (maximum 4.91 m). Extremely low Danube water level influenced the outflow of the water from the floodplain to the main channel, with a minimal water level of -0.30 m recorded in December 2011. In summer during 1973 and 2012 floodplain went through a second dry phase during chosen periods, lasting till the end of the studies.

Table 1. Danube water level (maximum, minimum and mean) and flooding dynamics of the river-floodplain system (categorization and duration) during the period of July 1972-October 1973 and July 2011-October 2012. (*) Approximation after Mikuska (1979) and Schwarz (2005)

Danube water level (WL, m)			1972-1973	2011-2012
Maximum			5.84	4.91
Minimum			0.55	-0.30
Mean			2.45	2.06
	Flooding categorization	Flooded area (%)*	Flooding duration (days/period)	
3.0-3.5	Minor	20	48	48
3.5-4.0	Moderate	40	43	38
4.0-5.0	Major	75	39	20
>5.0	Extremely high	>90	27	0
Total >3m			157	106
>3 in continuum			81	28




Figure 3. Daily courses of the Danube water level at 1,401.4 r. km during the period of July 1972-October 1973 (*dashed line*) and July 2011-October 2012 (*solid line*). Flooding categorization is shown in Table 1.

Table 2. Mean, minimum and maximum values of physical and chemical parameters in Lake Sakadaš during the period of July 1972-September 1973 and July 2011-October 2012.

Parameter (abbreviation, SI)	1972-1973			2011-2012		
	Mean	Min.	Max.	Mean	Min.	Max.
Water depth (WD, m)	7.8	6.7	9.9	6.5	5.4	7.7
Transparency (T, m)	0.9	0.2	1.8	1.0	0.7	1.9
Water temperature (WT, °C)	16.0	3.0	30.0	18.4	4.1	30.6
Dissolved oxygen (DO, mg L ⁻¹)	9.9	2.2	21.8	11.6	7.1	15.3
pH	8.2	7.7	9.7	8.3	7.7	9.0
Ammonium (NH ₄ ⁺ , µg L ⁻¹)				123.2	5.0	454.0
Nitrates (NO ₃ ⁻ , µg L ⁻¹)				861.6	20.0	3,950.0
Nitrites (NO ₂ ⁻ , µg L ⁻¹)				23.9	5.3	83.6
Organic nitrogen (orgN, µg L ⁻¹)				2,047.1	322.6	5,600.0
Total nitrogen (TN, µg L ⁻¹)				3,008.7	587.9	8,799.8
Total phosphorus (TP, µg L ⁻¹)				211.8	60.7	422.0
Conductivity (Cond, µS cm ⁻¹)				405.1	354.0	568.0
Chlorophyll <i>a</i> (Chl- <i>a</i> , µg L ⁻¹)				45.5	14.3	90.8
Chlorophyll <i>b</i> (Chl- <i>b</i> , µg L ⁻¹)				7.9	1.3	24.1
Chlorophyll <i>c</i> (Chl- <i>c</i> , µg L ⁻¹)				16.5	4.1	40.5

Variations of environmental parameters during the period of 1972-1973 (physical parameters) and 2011-2012 (physical and chemical parameters) are shown in Table 2. and Figures 4.-8. During the period 1972-1973, when Danube water level rose above 5 m and most of the floodplain area was flooded, the water depth of the lake reached 9.9 m. Lower water depth characterized the period of 2011-2012 with a maximum of 7.7 m. Water depth of the lake showed a significant positive correlation with Danube water level in 1972-1973 ($r = 0.78$, $p < 0.05$) and in 2011-2012 ($r = 0.73$, $p < 0.05$). During the summer months, transparency values were minimal in both periods, while in the winter months transparency reached maximal values. Water temperature was in opposite trending than transparency, which resulted in a significant negative correlation of parameters in both periods (1972-1973 ($r = -0.60$, $p < 0.05$) and in 2011-2012 ($r = -0.78$, $p < 0.05$)). Values of pH were higher than 7 during both periods indicating alkaline conditions. In the period of 1972-1973, pH and water temperature were in a significant positive correlation ($r = 0.64$, $p < 0.05$), while in 2011-2012 they were in a significant negative correlation ($r = -0.54$, $p < 0.05$). Higher mean value of dissolved oxygen was in 2011-2012, while in 1972-1973 values of dissolved oxygen were highly variable with a range from 2.2 to 21.8 mg L⁻¹.

Chlorophyll *a* in 2011-2012 oscillated between 14.3 to 90.8 $\mu\text{g L}^{-1}$ and was in a significant negative correlation ($r = -0.56$, $p < 0.05$) with transparency. Maximal values of Chlorophyll *a*, *b* and *c* were measured in October 2011, while minimum values were measured in January and May 2012. Conductivity (354-568 $\mu\text{S cm}^{-1}$) was in a significant negative correlation ($r = -0.62$, $p < 0.05$) with the water temperature.

Chemical parameters of Lake Sakadaš in 2011-2012 were highly variable. Highest values of ammonium (454 $\mu\text{g L}^{-1}$) and nitrates (3,950.0 $\mu\text{g L}^{-1}$) were noted in December 2011 during the highest transparency. Values of ammonium and nitrates showed a significant positive correlation (NH_4 ($r = 0.55$, $p < 0.05$); NO_3 ($r = 0.58$, $p < 0.05$)) with transparency. Concentrations of total nitrogen were higher in 2012 with a maximum of 8,799.8 $\mu\text{g L}^{-1}$. Values of total phosphorus had high oscillations (60.7-422 $\mu\text{g L}^{-1}$) during the whole period of 2011-2012.




Figure 4. Temporal variations of water depth and transparency in Lake Sakadaš during the periods of July 1972-September 1973 (*dashed line*) and July 2011-October 2012 (*solid line*).




Figure 5. Temporal variation of water temperature in Lake Sakadaš during the periods of July 1972-September 1973 (*dashed line*) and July 2011-October 2012 (*solid line*).




Figure 6. Temporal variation of dissolved oxygen in Lake Sakadaš during the periods of July 1972-September 1973 (*dashed line*) and July 2011-October 2012 (*solid line*).




Figure 7. Temporal variation of pH in Lake Sakadaš during the periods of July 1972-September 1973 (*dashed line*) and July 2011-October 2012 (*solid line*).




Figure 8. Temporal variations of ammonium (A), nitrates (B), nitrites (C), total nitrogen (D), total phosphorus (E), conductivity (F) and Chlorophyll *a* (G) in Lake Sakadaš during the period of July 2011-October 2012.

3.2. Taxonomical and functional diversity of phytoplankton

A total number of 317 phytoplankton taxa were registered during the periods in Lake Sakadaš. Chlorophyta and Bacillariophyceae achieved high species richness (Figure 9.). During the period 2011-2012, Cyanobacteria were represented by 33 taxa and differed from 1972-1973 with 23 taxa. In the period 1972-1973, there were 202 taxa identified with a range of 21 to 88 taxa per sample (Figure 10. (A)) with high oscillations of phytoplankton biomass ranging from 24.3 to 9,075.9 mg l⁻¹ in addition to the mean value of 995.0 mg l⁻¹. During the studied period 2011-2012, 190 taxa were identified with a range of 26 to 85 taxa (Figure 10. (B)) and lower values of biomass ranging from 3.1 to 146.2 mg l⁻¹ with the mean value of 34.5 mg l⁻¹. Species number of 1972-1973 had a significant positive correlation with water temperature ($r = 0.69$, $p < 0.05$) and pH ($r = 0.78$, $p < 0.05$), while biomass significantly correlated with transparency ($r = 0.59$, $p < 0.05$), dissolved oxygen ($r = 0.64$, $p < 0.05$) and pH ($r = 0.94$, $p < 0.05$). Number of species in 2011-2012 significantly correlated only with water temperature ($r = 0.60$, $p < 0.05$).





Figure 9. Percentage of phytoplankton taxonomical groups based on the total number of species in Lake Sakadaš during the periods of July 1972-September 1973 (A) and July 2011-October 2012 (B).

(A)



(B)




Figure 10. Temporal variations in phytoplankton total biomass and number of species in Lake Sakadaš during the periods of July 1972-September 1973 (A) and July 2011-October 2012 (B)

The identified phytoplankton taxa were sorted into 28 functional groups (FG): **A, B, C, D, E, F, G, H1, H2, J, K, L₀, M, MP, N, P, S1, S2, S_N, T, T_B, W1, W2, W_S, X1, X2, X3** and **Y**. Functional groups and representative species with their maximal contribution to total biomass within chosen periods are shown in Table 3.

The period of 1972-1973 was characterized with 27 FGs of which 16 FGs had relative biomass higher than 5% (Figure 11. (A)). Groups **B, D, G** and **S2** had the highest contribution to total biomass. Among 25 FGs from 2011-2012, 17 FGs had relative biomass higher than 5% (Figure 11. (B)) with best-developed groups **C, D** and **S1**. Comparison of dominant species within FGs in both periods had shown a significant difference in five groups: **B** (*Aulacoseira italica* → *Lindavia comta*), **C** (*Asterionela formosa* → *Cyclotella meneghiniana*), **H1** (*Anabaena planctonica* → *Cuspidothrix issatschenkoi*), **S1** (*Limnithrix redekei* → *Planktothrix agardhii*) and **S_N** (*Raphidiopsis mediterranea* → *Cylindrospermopsis raciborskii*). In 1972-1973 group **D, G, B** and **S2** had relative biomass higher than 70% of total biomass (Figure 12. (A)). In June 1972 group **D** reached 83.75% of total biomass with the high dominance of centric diatoms *Stephanodisucus hantzschii* and *Cyclostephanos dubius*. Group **G** reached up to 82.98% of total biomass in July 1973 with the dominance of *Eudorina elegans*, *Pleodorina illinoisensis* and *Pandorina morum*. Dominant species *Aulacoseira italica* from group **B** was massively developed in June 1973 with 77.88% of total biomass. Group **S2** with monodominant species *Spirulina* sp. reached up to 70.42% in September 1972.

During the studied period 2011-2012 only three functional groups developed with a contribution to total biomass higher than 40% (Figure 12. (B)). Highest contribution to total biomass was recorded in September 2011 during the massive development of filamentous Cyanobacteria from groups **S1, H1** and **S_N**, when only the **S1** reached up to 50.39% of total biomass with the dominance of *Planktothrix agardhii* (48.12%). Group **H1** reached maximal contribution to total biomass of this group during the studied period with 23.00% of *Dolichospermum sigmoideum* in August 2012. Filamentous Cyanobacteria functional groups were present during the whole studied period with higher contribution of **S1** group to the total biomass, except in June and July during the summer flood when phytoplankton composition was characterized with group **D**. *Ulnaria ulna* (31.48%) and *Stephanodisucus hantzschii* (4.43%) took dominant position during the whole period in group **D**. Group **C** with *Cyclotella meneghiniana* reached 48.69% of the total biomass in November 2011.

Group **Y** was continuously present during the periods with the highest contribution to total biomass in March 1973 (65.25%) and March 2012 (37.22%).

Besides mentioned FGs, significant difference in species composition within FGs comparing two periods was also found in groups **E, F, J, MP, T, W2** and **X2**. In May 2012 group **J** had a significant development with 24.08% of total biomass, while in the period 1973-1972, had a contribution to total biomass lower than 10%. Group **W2** was highly contributed with 45.68% of the total biomass in December 1972 with dominant species *Strombomonas annulata*.

Table 3. The maximal contribution of functional groups and species to total biomass in Lake Sakadaš during the period of July 1972-September 1973 and July 2011-October 2012.

Functional group	1972-1973		2011-2012	
	Maximal contribution of FG to total biomass (%)	Maximal contribution of species to total biomass (%)	Maximal contribution of FG to total biomass (%)	Maximal contribution of species to total biomass (%)
A	1.83	<i>Acanthoceras zachariasii</i> (Brun) Sim. (1.83%)	2.38	<i>Cyclotella</i> sp. (2.38%)
B	78.59	<i>Aulacoseira italica</i> (Ehrenb.) Sim. (77.88%)	1.39	<i>Lindavia comta</i> (Kütz.) Nakov, Gullory, Julius, Theriot & Alverson (1.39%)
C	0.60	<i>Asterionella formosa</i> Hass. (0.60%)	48.69	<i>Cyclotella meneghiniana</i> Kütz. (48.69%)
D	83.75	<i>Ulnaria ulna</i> (Nitz.) Comp. (66.45%); <i>Stephanodiscus hantzschii</i> Grun. (41.90%); <i>Cyclostephanos dubius</i> (Hust.) Round (41.28%)	40.73	<i>U. ulna</i> (35.59%); <i>S. hantzschii</i> (25.64%); <i>Ulnaria acus</i> (Kütz.) Aboal (6.19%)
E	13.08	<i>Dinobryon divergens</i> var. <i>angulatum</i> (Seligo) Brunnthaler (10.68%)	1.87	<i>Dinobryon divergens</i> Imh. (1.87%)
F	7.29	<i>Oocystis marssonii</i> Lemm. (6.78%)	5.35	<i>Micractinium bornhemienae</i> (W.Conrad) Korshikov (3.74%)
G	82.84	<i>Eudorina elegans</i> Ehrenb. (29.79%); <i>Pleodorina illinoisensis</i> Kofoid (27.22%); <i>Pandorina morum</i> (O.F.Müller) Bory (25.83%)		
H1	3.89	<i>Anabaena planctonica</i> Brunnthaler (2.00%)	24.38	<i>Cuspidothrix issatschenkoi</i> (Usachev) P.Rajaniemi, Kom., R.Willame, P.Hrouzek, K.Kastovská, L.Hoffm. & K.Sivonen (21.79%); <i>Dolichospermum sigmoides</i> (Nygaard) Wacklin, L.Hoffm. & Kom. (9.54%); <i>Aphanizomenon flosaquae</i> Ralfs ex Born. & Flah. (6.80%); <i>Dolichospermum solitarium</i> (Kleb.) Wacklin, L.Hoffm. & Kom. (5.94%)
H2	0.53	<i>Gloeotrichia</i> sp. (0.53%)		
J	9.55	<i>Pediastrum boryanum</i> var. <i>boryanum</i> (Turp.) Menegh. (6.63%)	24.09	<i>Tetradismus lagerheimii</i> M.J.Wynne & Guiry (11.34%); <i>Coelastrum microporum</i> Nägeli (5.21%)
K	0.04	<i>Aphanothece elabens</i> (Bréb. ex Menegh.) Elenkin (0.04%)	3.64	<i>Aphanocapsa delicatissima</i> West & G.S.West (3.64%)
L ₀	6.63	<i>Peridinium cinctum</i> (O.F.Müller) Ehrenb. (4.45%)	10.38	<i>Peridinium aciculiferum</i> Lemm. (6.75%); <i>P. cinctum</i> (5.29%)
M	0.64	<i>Microcystis aeruginosa</i> (Kütz.) Kütz. (0.32%); <i>Microcystis wesenbergii</i> (Kom.) Kom. ex Kom. (0.32%)	6.83	<i>M. aeruginosa</i> (6.59%)
MP	17.50	<i>Brachysira exilis</i> (Kütz.) Round & D.G.Mann (11.52%); <i>Gyrosigma macrum</i> (W.Smith) J.W.Griffith & Henfrey (7.56%); <i>Oscillatoria tenuis</i> C.Agardh ex Gomont (5.92%)	6.12	<i>Amphora ovalis</i> (Kütz.) Kütz. (5.57%)
N	0.05	<i>Cosmarium</i> sp. (0.03%)	3.24	<i>Cosmarium phaseolus</i> Bréb. ex Ralfs (3.24%)
P	10.79	<i>Staurastrum</i> sp. (10.37%); <i>Closterium macilentum</i> Bréb. (8.58%); <i>Aulacoseira granulata</i> (Ehrenb.) Sim. (6.56%)	11.82	<i>A. granulata</i> (7.89%)
S1	1.43	<i>Limnithrix redekei</i> (Goor) Meffert (1.38%)	50.39	<i>Planktothrix agardhii</i> (Gom.) Anag. & Kom. (48.12%); <i>L. redekei</i> (14.93%); <i>Pseudanabaena limnetica</i> (Lemm.) Kom. (9.97%)
S2	70.42	<i>Spirulina</i> sp. (70.42%)		
S _N	10.08	<i>Raphidiopsis mediterranea</i> Skuja (10.08%)	5.85	<i>Cylindrospermopsis raciborskii</i> (Wol.) Seenayya & Subba Raju (5.85%)
T	17.03	<i>Mougeotia</i> spp. (17.03%)	9.12	<i>Binuclearia lauterbornii</i> (Schmidle) Proschkina-Lavrenko (7.98%)
T _B	3.55	<i>Navicula rhynchocephala</i> Kütz. (1.69%)	1.16	<i>Navicula</i> sp. (0.81%)
W1	14.27	<i>Lepocinclis ovum</i> (Ehrenb.) Lemm. (13.09%)	7.90	<i>L. ovum</i> (6.97%); <i>Euglena texta</i> (Duj.) Hübner (6.77%)
W2	45.68	<i>Strombomonas annulata</i> (Daday) Deflandre (44.21%); <i>Trachelomonas volvocina</i> (Ehrenb.) Ehrenb. (5.93%); <i>Trachelomonas hispida</i> var. <i>Crenulocollis</i> (Maskell) Lemm. (5.51%)	4.13	<i>T. volvocina</i> (1.65%)
W _S	6.84	<i>Synura uvella</i> Ehrenb. (6.84%)	6.76	<i>S. uvella</i> (6.76%)
X1	0.84	<i>Pseudoditymocystis planctonica</i> (Korshikov) E.Hegewald & Deason (0.82%)	4.58	<i>P. planctonica</i> (3.45%)
X2	0.45	<i>Chlamydomonas globosa</i> J.W.Snow (0.45%)	16.57	<i>Carteria</i> sp. (8.50%); <i>Rhodomonas</i> sp. (7.63%); <i>Rhodomonas lacustris</i> Pasch. & Rutt. (6.81%); <i>Chlamydomonas</i> sp. (6.09%)
X3			5.50	<i>Chrysococcus rufescens</i> Klebs (3.69%)
Y	65.25	<i>Cryptomonas erosa</i> Ehrenb. (62.08%); <i>Naiadinitium polonicum</i> (Wolosz.) S.Carty (46.93%)	37.22	<i>Cryptomonas ovate</i> Ehrenb. (27.87%); <i>Cryptomonas</i> sp. (21.76%); <i>C. erosa</i> (13.43%)




Figure 11. Dynamics of dominant functional groups on relative biomass in Lake Sakadaš during the period of July 1972-September 1973 (A) and July 2011-October 2012 (B). FGs with relative biomass higher than 5%.





Figure 12. The maximal contribution to total biomass of best developed functional groups in Lake Sakadaš during the period of July 1972-September 1973 (A) and July 2011-October 2012 (B).




Figure 13. Non-metric multidimensional scaling (nMDS) plot based on biomass of phytoplankton functional groups in Lake Sakadaš during the periods July 1972-September 1973 and July 2011-October 2012. Circles indicate samples from 1972-1973, while triangles indicate samples from 2011-2012. *Solid lines* separate groups, *dashed lines* separate subgroups.

The multivariate nMDS analysis displayed a clear separation of samples into three groups based on the biomass of phytoplankton FGs in the lake during 1972-1973 and 2011-2012 (Figure 13.). Group 2011-2012 encompassed samples from the period of 2011-2012 with the division in three subgroups characterized with Bacillariophyceae, Cryptophyceae and Cyanophyceae species. Lower biomass characterized subgroup with the dominance of **Y**, **D**, **S1** and **X2** groups including samples from December 2011-April 2012 with the highest contribution of group **Y**. The subgroup with the dominance of **Y**, **J**, **D** and **S1** included samples from May-July and September-October 2012, while subgroup with the dominance of **S1**, **H1**, **Y** and **D** groups included samples from July-October 2011 and August 2012.

Group 1972-1973 encompassed subgroup with samples from December 1972-February 1973 and May 1973 with high contribution to total biomass of **D** and **W2** group, and the subgroup with the dominance of **Y** and **D** (September-November 1972 and March-April 1973).

Group 1972-1973 with monodominant FGs represented samples with the dominance of one FG whose contribution was higher than 70% of the total biomass.

Samples from November 2011 and August 1972 had high dominance of **D** group and were not included in any of current subgroups, but they were listed in current groups while they had more similarity with samples from the group than with one another.

3.3. Phytoplankton assemblage index

Factor numbers were determined for each functional group identified in Lake Sakadaš from the period of July 1972-September 1973 and July 2011-October 2012. Determined *F* values were presented in Table 4. with *F* values of similar water body types.

Q index values of the lake in period 1972-1973 varied from 0.17 to 3.10 (Figure 14. (A)). The ecological status of the lake was under the *medium* classification during the most of the period, except in July and September 1973 when the lake was under the *bad* classification, December 1972 and January in *tolerable*, and in May 1973 when it was in *good* condition.

Q index values in the period 2011-2012 varied with season, changing the ecological status of Lake Sakaš from *tolerable* to *good* (Figure 14. (B)). The highest qualification with 3.68 occurred in November 2011, and the lowest evaluation with 1.04 in September 2011 when the cyanobacterial bloom occurred.

Mean values of Q index in chosen periods showed that Lake Sakadaš has *medium* ecological status.

Table 4. Factor number (F) of phytoplankton functional group of Lake Sakadaš with comparison to F for similar water body types: oxbow outside flood control dams (Hungary)-Padisák et al. (2006); shallow eutrophic reservoir (Brazil)-Crossetti and Bicudo (2008), water-supply reservoir (Brazil)-Becker et al. (2010); small and middle-sized lowland lakes (Bulgaria)-Belkinova et al. (2014).

Functional groups	Factor numbers	F (Padisák et al., 2006)	F (Crossetti and Bicudo, 2008)	F (Becker et al., 2010)	F (Belkinova et al., 2014)
A	5	5			
B	2	5			5
C	5	5		3	5
D	2	3	2	2	2.5
E	5	5	5	5	5
F	3	3	5	2	5
G	0	4			3
H1	1	1	1	0	1
H2	3	3			
J	5	5	5	2	3
K	2	2	3	4	3
L ₀	5	5	5	4	5
M	0	0	0		
MP	3	3		1	3
N	5	5	5	5	5
P	5	5	2	0	5
S1	0	0	0	0	0
S2	0	2			0
S _N	0	0	0		
T	5	5	5		5
T _B	5				
W1	2	2	0	0	
W2	0	3	1	1	
W _S	3	4			
X1	3	3	5	3.5	2.5
X2	3.5	3.5	5	5	3
X3	4	4	4	5	5
Y	3	3.5	3	3	3.5




Figure 14. Dynamics of Q index and ecological status evaluation of Lake Sakadaš during the period of July 1972-September 1973 (A) and July 2011-October 2012 (B).

4. DISCUSSION

A comparison of historical records (1972-1973) with recent data (2011-2012) showed important differences in the abundance and dynamics of the dominant phytoplankton assemblages in the investigated floodplain lake.

As a consequence of strong anthropogenic influence in the past, the lake environment was dramatically deteriorated by the inflow of wastewaters from surrounding agricultural area. Large amounts of suspended matter supported the intensive decomposition process and resulted in the periodical appearance of the hypoxic condition and dramatic increase of pH. After four decades of water protection in the whole floodplain area and applied revitalization measure (sediment removal) in the lake, it seems that the lake has recovered. Investigations done in the past few decades showed that flooding dynamic, still preserved in its near-natural state, is the key ecological factor that influenced phytoplankton dynamics (Mihaljević et al., 2009). Moreover, the results of the effectiveness of the functional approach designed to explain the phytoplankton changes associated with hydrological events showed that the functional classification allows representing of the hydrological phases (Stević et al., 2013). The instrument applied to assess the changes in the ecological status of this floodplain lake, Q index, was applied because unlike other indices it can reflect general anthropogenic pressure (Padisák et al., 2006). Due to the known fact that determination of factor number (F) for each FG is the most sensitive step in defining of Q index (Belkinova et al., 2014), a comparison between FGs in impacted and semi-natural conditions allows to allocated lower values to FGs found to be the most expressive in the undesirable conditions.

Given the dataset of phytoplankton functional groups and the main group arrangements in the multivariate nMDS analysis showed variation related to the past and recent condition. Heavy algal bloom (with extremely high biomass) of species assemblages from one FGs or bloom of only one species from particular FGs were the most distinctive feature of the environmental conditions in the past. The bloom of species from the group **G** (*Eudorina*, *Pandorina*, *Pleodorina*), when the relative biomass reached up to 80% of the total biomass, appeared in July 1973 in the highly alkaline conditions (pH 9.7). As was noted by Gucunski (1975), the lake water at that time had an intensively green color and “there was no fish in the lake”. During the summer of 1973, Kiss (1977) also recorded intensively algal bloom of *Eudorina elegans* in the Tisza dead-arm in strongly polluted salt and alkaline waters, containing large amounts of decomposing organic matters and qualified those ecological state as “eutrophication with a

dangerous extent”. Heavy polluted water has been widely recognized as a suitable environment for blooming of species from the group **G**. Almost monodominant *Eudorina* blooms were registered in the polluted shallow pond which receives large quantities of sewage from the nearby city (Munawar and Zafar, 1967). Immense bloom of *Pandorina morum* occurred in the Lake Erie (Laurentian Great Lakes), which has a long history of anthropogenic perturbation with excessive nutrient loading (Millie et al., 2009). According to Reynolds et al. (2002), habitat template for group **G** is characterized by nutrient-rich conditions in stagnating water columns, small eutrophic lakes and very stable phases in larger river-fed basins and storage reservoirs. Different weights of factor *F* were used in the qualification of the **G** in order to determine ecological status: value 4 for oxbow lakes outside flood control, value 0 for different types of alkaline lakes in Hungary (Padisák et al., 2006), value 3 for small and middle-size lowland lakes in Bulgaria (Belkinova et al., 2014) and value 1 for the reservoir in China (Wang et al., 2011). In this study, factor *F* with value 0 is used and calculated *Q* index for July 1973 is 0.17, reflecting the *bad* ecological status of the lake, which was affected by wastewaters and was in strongly alkaline condition.

Another heavy algal bloom occurred thereafter (September 1973), formed by filamentous blue-green alga *Spirulina* sp. (group **S2**), which was contributed more than 70% to the total biomass, confirmed heavy polluted conditions in the lake. It is known that *Spirulina* bloom occurs widely in polluted lakes, particularly in tropical and sub-tropical lakes with high salinity (El-Bestawy et al., 1996). Warm, shallow and often highly alkaline waters are habitat template for the group **G** (Reynolds et al., 2002) and it is clear that used factor *F* value for the group **S2** is 0. Moreover, current research showed that the group **S2** can be associated with other shade tolerant cyanobacterial species from **S_N** assemblage, forming an **S2-S_N** association in shallow polymictic tropical lakes (Gebrehiwot et al., 2017). Accompanied species from **S_N** group during the bloom of *Spirulina* in the Lake Sakadaš was *Raphidiopsis mediterranea*. The occurrence of this subtropical cyanobacterial species was registered for the first time during summer 1973 in all floodplain habitats of Kopački Rit, and also in the main riverbed of the Danube (Gucunski, 1975). The first appearance of *Raphidiopsis mediterranea* in the Lake Balaton was also noted in summer of 1973 (Padisák, 1992). However, after these findings there are no records of *R. mediterranea* in waters of Kopački Rit, in spite that distribution pattern of this species showed its continually spreading throughout various European freshwater ecosystems from the beginning of this century (Kaštovský et al., 2010). It should be emphasized that the morphology of *Raphidiopsis* is highly similar to that of *Cylindrospermopsis*, especially in the non-

heterocytous life-cycle stages of *Cylindrospermopsis* species where heterocystes and kinetes are not formed (Li et al., 2001; Moustaka-Gouni et al., 2009; Alster et al., 2010). According to Moustaka-Gouni et al. (2009), *R. mediterranea* can represent non-heterocytous stages of *Cylindrospermopsis* complex life cycle. Thus, overlapping features of *Cylindrospermopsis/Raphidiopsis* have led to the difficulty in distinguishing these two genera by the traditional taxonomic approach.

A great expansion of *C. raciborskii* in the temperate waters was noted during the past decades (Padisák, 1997). After its first records in the Danube Delta (Roll, 1961), its mass development was found in various floodplain habitats along the Danube River, from the Lower Danube (Stoyneva, 2003) till the Upper Danube (Dokulil and Mayer, 1996). The first finding of *C. raciborskii* bloom in the Lake Sakadaš was in the extremely dry summer of 2003 when its long-lasting over-domination led to the establishment of equilibrium phase (Mihaljević et al., 2009). Afterwards, massive development of *C. raciborskii* has been noted in the summer of 2007 (Mihaljević and Stević, 2011). Given data showed that *C. raciborskii* was accompanied species (5.85 % of the total biomass) during the blooming of cyanobacteria, dominated by species from the **S1** group (46.87% of the total biomass, over-dominated by *Planktothrix agardhii*) and co-dominated by species from **H1** group (24.38 % of the total biomass). It is known that species belonging to the **S1**, **H1** and **S_N** group are frequently in competition (Padisák et al., 2003). A comprehensive research of cyanobacterial blooming in this floodplain lake showed that low-nitrogen **H1** group was particularly sensitive to stress caused by flooding, while filamentous N₂-fixing (**S_N**) and non-N₂-fixing species (**S1**) showed tolerance to short-term flooding (Stević et al., 2013). Thus, in the highly turbulent environmental conditions in the floodplain lake, species of the **H1** group are less successful in forming long-lasting bloom. Factor *F* value 1 is used for the group **H1**, while for the groups **S1**, **S2**, **S_N** factor *F* is 0.

Concerning the appearance of cyanobacterial bloom in the investigated lake it must be emphasized that in the past decades, as a consequences of frequent occurrence of extreme flood events, the lake can be shifted between a state of turbid water, characterized by high phytoplankton biomass and regular appearance of cyanobacteria blooms, to a state of clear water with very low phytoplankton biomass and absence of cyanobacteria, and back to the turbid state (Mihaljević and Stević, 2011). It is evident that the appearance of cyanobacterial bloom has a strong influence on the ecological status of the lake and values of Q index were inversely proportional to the dominance of cyanobacterial groups. Thus, low values of Q index

(1.04 and 1.54) during the cyanobacterial bloom in September-October 2011 substantially decreased the final assemblage index value on an annual scale (2.57).

A permanent component of phytoplankton in the lake are diatoms, sharing a certain percentage on a whole-year scale in both observed periods. A previous research (Stević et al., 2013) summarized that flooding of the lake and mixed waters are favorable for the development of diatoms belonging to the **B**, **C**, **D** and **P** functional groups. However, due to the dilution and washout effect, their biomass is low during the long-lasting flooding despite their input from the river. The similar pattern was established in the investigated period 2011-2012. However, comparison with the historical records showed that in particular conditions some diatoms species can be mass developed and form a heavy bloom. The most expressive diatom bloom had occurred in June of 1973, after extremely high flooding and in the hypoxic conditions, when species from the **D** group (*Stephanodiscus hantzschii* and *S. dubius*) contributed up to 83.75% of the total biomass. Similar physical and chemical characterization of the environment, hypoxic and alkaline condition after extremely high flooding in July of 1972, supported a heavy monotypic bloom of centric diatom *Aulacoseira italica* from the group **B**, which contributed up to 78.59% of the total biomass. The high abundance of *Stephanodiscus hantzschii* is considered as indicative of high nutrient levels and is a good indicator of eutrophic conditions in lakes (Tolotti et al., 2010, and cites therein). It is known that many species of *Stephanodiscus* and *Aulacoseira* form a vegetative resting cell in surficial lake sediments and are meroplanktonic, spending a part of their life in hypolimnion or the littoral zone and found in the plankton only during periods when the water column undergoes turbulent mixing (Lashaway and Carrick, 2010). Moreover, Yang et al. (2015) revised the physiological characteristics defined by Reynolds et al. (2002) for the groups **B** and **D** as follows. Habitat template for the group **B** (*Aulacoseira italica*): mixed, mesotrophic, small-medium lakes; adapted to low light; some species could form resting cells, meroplanktonic. Habitat template for the group **D** (*Stephanodiscus hantzschii*): shallow, nutrient-enriched, well-ventilated waters; sensitive to nutrient depletion; some species could form resting cells. Factor *F* value used till now for more or less similar lake types (Padisák et al., 2006; Belkinova et al., 2014; Wang et al., 2011) varied between 1 and 5 for the group **B** and between 2 and 3 for the group **D**. According to the given experience of heavy blooming of species from **B** and **D** in the condition of extremely high anthropogenic pressure and deteriorated waters factor *F* value 2 was used for both groups.

The appearance of diatom-dominated functional groups **C**, **P**, **Tb** and **A**, groups with maximal *F* factor value 5, is commonly associated with the conditions of flooding phase and clear water

with very low phytoplankton biomass (Mihaljević and Stević, 2011; Stević et al., 2013), as confirmed by the obtained results. Species from the group **T_B** (*Navicula* spp., *Cymbella*, *Melosira varians*) characteristic for highly lotic environments (Padisák et al., 2009; Abonyi et al., 2012) as well as from the group **A** (small centric diatoms) characteristic for clear, often mixed lakes with low P half-saturation (Reynolds et al., 2002) occasionally occurred with very small contribution (less than 5%) to the total biomass. Likewise, species from the group **P** (*Aulacoseira granulata*) which is characteristic for more eutrophic waters (Yang et al., 2016). It should be noted that the dependence of species from the group **P** upon physical mixing is strongly apparent, requiring a continuous or semi-continuous mixed layer of 2-3 m in thickness (Reynolds et al., 2002). Mixing of the water in the floodplain lake is related to the flood pulses, which can support re-suspension of meroplanktonic species, such as *A. granulata*, in the water column. However, flooding usually lasts a short time and is followed with dry phase with stagnant water unfavorable for further development of species from the group **P**. However, small centric diatoms from the group **C** were permanently present in phytoplankton during the period 2011-2012, while their occurrence in the past was insignificant. Their dominance (48.69% of the total biomass) in November 2012 coincided with the decrease of nutrients in the lake water and Q index reached a maximum value of 3.68, which indicated *good* ecological status of the lake in that time.

5. CONCLUSION

A comparison of previous and ongoing assessments of phytoplankton assemblage dynamics has greatly assisted in the evaluation of the ecological state of the floodplain lake using the assemblages Q index, especially because there is no data for the determination of pristine condition. The chosen historical assessment reflects a period of significant anthropogenic influences that resulted in rapid eutrophication, moreover deterioration of water environment. The frequent appearance of heavy bloom of only one species or assemblages characteristic for polluted lakes decreased the value of Q index to *bad* and *tolerable* state. Recent data suggest that water quality improvement and near-natural hydrological condition support algal assemblages characteristic for naturally eutrophic lakes and values of Q index varied between *medium* to *good* ecological status. Currently, the most important threat for this lake is a long-lasting bloom of alien invasive cyanobacteria what might lead to shifting towards the *bad* condition. Altogether, by the evaluation of historical records and the application of assemblages Q index, natural changes can be distinguished from anthropogenic changes in the floodplain lake.

6. REFERENCES

- Abonyi A, Leitão M, Lançon AM, Padisák J. 2012. Phytoplankton functional groups as indicators of human impacts along the River Loire (France). *Hydrobiologia* 698 (1):233-249.
- Alster A, Kaplan-Levy RN, Sukenik A, Zohary T. 2010. Morphology and phylogeny of a non-toxic invasive *Cylindrospermopsis raciborskii* from a Mediterranean Lake. *Hydrobiologia* 639:115-128.
- Amoros C, Bornette G. 2002. Connectivity and biocomplexity in waterbodies of riverine floodplains. *Freshwater Biology* 47:761:776.
- Anagnostidis K, Komárek J. 1985. Modern approach to the classification system of cyanophytes. 1. Introduction. *Archiv für Hydrobiologie* (Supplement) 71:291-302.
- Anagnostidis K, Komárek J. 1988. Modern approach to the classification system of cyanophytes. 3. Oscillatoriales. *Archiv für Hydrobiologie* (Supplement) 80:327-472.
- APHA (American Public Health Association). 1992. *Standard methods for the examination of water and wastewater*. Washington, DC: American Public Health Association.
- Barone R, Castelli G, Naselli-Flores L. 2010. Red sky at night cyanobacteria delight: the role of climate in structuring phytoplankton assemblage in a shallow, Mediterranean lake (Biviere di Gela, southeastern Sicily). *Hydrobiologia* 639:43-53.
- Becker V, Caputo L, Ordóñez J, Marcé R, Armengol J, Crossetti LO, Huszar VLM. 2010. Driving factors of the phytoplankton functional groups in a deep Mediterranean reservoir. *Water Research* 44:3345-3354.
- Becker V, de Souza Cardoso L, Huszar VLM. 2009a. Diel variation of phytoplankton functional groups in a subtropical reservoir in southern Brazil during an autumnal stratification period. *Aquatic Ecology* 43:285-293.
- Becker V, Huszar VCM, Crossetti LO. 2009b. Responses of phytoplankton functional groups to the mixing regime in a deep subtropical reservoir. *Hydrobiologia* 628:137-151.
- Belkinova D, Padisák J, Gecheva G, Cheshmedjiev S. 2014. Phytoplankton based assessment of ecological status of Bulgarian lakes and comparison of metrics within the Water Framework Directive. *Applied Ecology and Environmental Research* 12(1):83-103.

- Bonacci O, Tadić Z, Moržan A, Radeljak I. 2002. *Nature Park Kopački Rit Management plan. Sector study: Hydrology and meteorology*. Zagreb: Elektroprojekt.
- Borges PAF, Train S, Rodrigues LC. 2008. Spatial and temporal variation of phytoplankton in two subtropical Brazilian reservoirs. *Hydrobiologia* 607: 63-74.
- Borics G, Tóthmérész B, Lukács BA, Várbíró G. 2012. Functional groups of phytoplankton shaping diversity of shallow lake ecosystems. *Hydrobiologia* 698:251-262.
- Borics G, Várbíró G, Grigorszky I, Krasznai E, Szabó S, Kiss KT. 2007. A new evaluation technique of potamoplankton for the assessment the ecological status of rivers. *Archiv für Hydrobiologie, Supplement Large Rivers* 161:465-486.
- Buijse AD, Coops H, Staras M, Jans LH, van Geest GJ, Grift RE, Ibelings BW, Oosterberg W, Roozen FCJM. 2002. Restoration strategies for river floodplains along large lowland rivers in Europe. *Freshwater Biology* 47:889-907.
- Çelekli A, Öztürk B. 2014. Determination of ecological status and ecological preferences of phytoplankton using multivariate approach in a Mediterranean reservoir. *Hydrobiologia* 740:115-135.
- Clarke KR, Warwick RM. 2001. *Change in marine communities: An approach to statistical analysis and interpretation* (2nd ed.). PRIMER-E: Plymouth.
- Costa LS, Huszar VLM, Ovalle AR. 2009. Phytoplankton functional groups in a tropical estuary: hydrological control and nutrient limitation. *Estuaries and Coasts* 32:508-521.
- Crossetti LO, Becker V, Cardoso LS, Rodrigues LR, Costa LS, Motta-Marques D. 2013. Is phytoplankton functional classification a suitable tool to investigate spatial heterogeneity in a subtropical shallow lake? *Limnologia* 43:157-163.
- Crossetti LO, Bicudo CEM. 2008. Phytoplankton as a monitoring tool in a tropical urban shallow reservoir (Garças Pond): the assemblage index application. *Hydrobiologia* 610:161-173.
- Dawidek J, Ferencz B. 2012. Hydrological processes in the riverine systems, the origin and classifications of floodplain lakes. *Ekológia (Bratislava)* 31(3):331-340.

- Demir AN, Fakioğlu Ö, Dural B. 2014. Phytoplankton functional groups provide a quality assessment method by the Q assemblage index in Lake Mogan (Turkey). *Turkish Journal of Botany* 38:169-179.
- Dokulil MT, Mayer J. 1996. Population dynamics and photosynthetic rates of a *Cylindrospermopsis* - *Limnothrix* association in a highly eutrophic urban lake, Alte Donau, Vienna, Austria. *Algological Studies* 83:179-195.
- El-Bestawy E, Bellinger EG, Sigeo DC. 1996. Elemental composition of phytoplankton in a subtropical lake: X-ray microanalytical studied on the dominant algae *Spirulina platensis* (Cyanophyta) and *Cyclotella meneghiniana* (Bacillariophyceae). *European Journal of Phycology* 31:157-166.
- Fonseca BM, Bicudo CEM. 2008. Phytoplankton seasonal variation in a shallow stratified eutrophic reservoir (Garças Pond, Brazil). *Hydrobiologia* 600:267-282.
- Gebrehiwot M, Demeke K, Stiers I, Triest L. 2017. Phytoplankton functional dynamics in a shallow polymictic tropical lake: the influence of emergent macrophytes. *Hydrobiologia* 797:69-86.
- Gucunski D, Popović Ž. 1984. Podaci za kvantitativnu analizu fitoplanktona rezervata Kopački rit. [Data for quantitative analysis of phytoplankton of the reservation Kopački Rit.] *ANALI zavoda Jugoslav. akad.* 2:277-298.
- Gucunski D. 1975. Kvantitativna istraživanja fitoplanktona u Upravljanom prirodnom rezervatu Kopački rit. Dissertation, University of Zagreb.
- Guiry MD, Guiry GM. 2014. AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. <http://www.algaebase.org>; searched in May 2017.
- Hajnal É, Padisák J. 2008. Analysis of long-term ecological status of Lake Balaton based on the ALMOBAL phytoplankton database. *Hydrobiologia* 599:227-237.
- Hindák F, Cyrus Z, Marvan P, Javornický P, Komárek J, Ettl H, Rosa K, Sladečková A, Popovsky J, Punčocharova M, Lhotsky O. 1978. *Slatkovodne riasy*. Slovenske pedagogicke nakladatelstvo, Bratislava.
- Horvatić J, Mihaljević M, Stević F. 2003. Algal growth potential of *Chlorella kessleri* Fott et Nov. In comparison with *in situ* microphytoplankton dynamics in the water of Lake Sakadaš marshes. *Periodicum Biologorum* 105:307-312.

- Hu R, Han B, Naselli-Flores L. 2013. Comparing biological classifications of freshwater phytoplankton: a case study from South China. *Hydrobiologia* 701:219-233.
- Hustedt F. 1976. *Bacillariophyta*. Otto Koeltz Science Publishers, Koenigstein.
- Huszar V, Kruk C, Caraco N. 2003. Steady state of phytoplankton assemblage of phytoplankton in four temperate lakes (NE USA). *Hydrobiologia* 502:97-109.
- Izaguirre I, Allende L, Escaray R, Bustingorry J, Pérez G, Tell G. 2012. Comparison of morpho-functional phytoplankton classifications in human-impacted shallow lakes with different stable states. *Hydrobiologia* 698 (1):203-216.
- Javornický P, Komárková J. 1973. The changes in several parameters of plankton primary productivity in Slapy Reservoir 1960-1967, their mutual correlations and correlations with the main ecological factors. In *Hydrobiological studies*, Hrbáček J, Straškraba M (eds). Academia: Prague, 155-211 pp.
- Kaštovský J, Hauer T, Mareš J, Krautová M, Bešta T, Komárek J, Desortová B, Heteša J, Hindáková A, Houk V, Janeček E, Kopp R, Marvan P, Pumann P, Skácelová O, Zapomělová E. 2010. A review of the alien and expansive species of freshwater cyanobacteria and algae in the Czech Republic. *Biological Invasions* 12:3599-3625.
- Kiss I. 1977. Investigation of the water blooms of *Eudorina elegans* in the dead-arm of the River Tisza at the community Mártély. *Tiscia (Szeged)* 12:37-47.
- Komárek J, Anagnostidis K. 1989. Modern approach to the classification system of cyanophytes. 4. Nostocales. *Algological Studies* 56:247-345.
- Kruk C, Huszar VLM, Peeters ETHM, Bonilla S, Costa L, Lüring M, Reynolds CS, Scheffer M. 2010. A morphological classification capturing functional variation in phytoplankton. *Freshwater Biology* 55 (3):614-627.
- Kruk C, Mazzeo N, Lacerot G, Reynolds CS. 2002. Classification schemes for phytoplankton: a local validation of a functional approach to the analysis of species temporal replacement. *Journal of Plankton Research* 24 (9):901-912.
- Kruk C, Segura AM. 2012. The habitat template of phytoplankton morphology-based functional groups. *Hydrobiologia* 698 (1):191-202.

Lashaway AR, Carrick HJ. 2010. Effects of light, temperature and habitat quality on meroplanktonic diatom rejuvenation in Lake Erie: implications for seasonal hypoxia. *Journal of Plankton Research* 32 (4):479-490.

Li R, Carmichael WW, Brittain S, Eaglesham GK, Shaw GR, Mahakhan A, Noparatnaraporn N, Yongmanitchai W, Kaya K, Watanabe MM. 2001. Isolation and identification of the cyanotoxin cylindrospermopsin and deoxy-cylindrospermopsin from a Thailand strain of *Cylindrospermopsis raciborskii* (Cyanobacteria). *Toxicon* 39:973-980.

Meffert ME, Oberhäuser R, Overbeck J. 1981. Morphology and Taxonomy of *Oscillatoria redekei* (Cyanophyta). *British Phycological Journal* 16:107-114.

Mieleitner J, Borsuk M, Bürgi HR, Reichert. 2008. Identifying functional groups of phytoplankton using data from three lakes of different trophic state. *Aquatic Sciences* 70:30-46.

Mihaljević M, Getz D, Tadić Z, Živanović B, Gucunski D, Topić J, Kalinović I, Mikuska J. 1999. *Kopački Rit-research survey and bibliography*. Croatian Academy of Arts and Sciences, Zagreb.

Mihaljević M, Špoljarić D, Stević F, Žuna Pfeiffer 2013. Assessment of flood-induced changes of phytoplankton along a river-floodplain system using the morpho-functional approach. *Environmental Monitoring and Assessment* 185:8601-8619.

Mihaljević M, Stević F, Horvatić J. 2009. Dual impact of the flood pulses on the phytoplankton assemblages in a Danubian floodplain lake (Kopački Rit Nature Park, Croatia). *Hydrobiologia* 618:77-88.

Mihaljević M, Stević F. 2011. Cyanobacterial blooms in a temperate river-floodplain ecosystem: the importance of hydrological extremes. *Aquatic Ecology* 45:335-349.

Mikuska J. 1979. Ekološke osobine i zaštita specijalnog zoološkog rezervata "Kopački rit" s posebnim osvrtom na ekologiju kraljevnjaka. Dissertation, University of Zagreb.

Millie DF, Fahnenstiel GL, Dyble Bressie J, Pigg RJ, Rediske RR, Klarer DM, Tester PA, Litaker RW. 2009. Late-summer phytoplankton in western Lake Erie (Laurentian Great Lakes): bloom distributions, toxicity, and environmental influences. *Aquatic Ecology* 43 (4):915-934.

Molina-Navarro E, Martínez-Perez S, Sastre-Merlín A, Verdugo-Althöfer M, Padisák J. 2014. Phytoplankton and suitability of derived metrics for assessing the ecological status in a limno-

reservoir, a Water Framework Directive nondefined type of Mediterranean waterbody. *Lake and Reservoir Management* 30 (1):46-62.

Moustaka-Gouni M, Kormas AK, Vardaka E, Katsiapi M, Gkelis S. 2009. *Raphidiopsis mediterranea* Skuja represents non-heterocytous life-cycle stages of *Cylindrospermopsis raciborskii* (Woloszynska) Seenayya et Subba Raju in Lake Kastoria (Greece), its type locality: Evidence by morphological and phylogenetic analysis. *Harmful Algae* 8:864-872.

Munawar M, Zafar AR. 1967. A preliminary study of vertical movement of *Eudorina elegans* and *Trinema lineare* during a bloom caused by them. *Hydrobiologia* 29 (1-2):140-148.

Naselli-Flores L, Padisák J, Dokulil MT, Chorus I. 2003. Equilibrium/steady-state concept in phytoplankton ecology. *Hydrobiologia* 502 (1-3):395-403.

Ochocka A, Pasztaleniec A. 2016. Sensitivity of plankton indices to lake trophic conditions. *Environmental Monitoring and Assessment* 188:622.

OECD. 1982. Eutrophication of waters. Monitoring, Assessment and Control. OECD Publications, Paris.

Ongun Sevindik T, Tunca H, Gönülol A, Yildirim Gürsoy N, Küçükkaya SN, Durgut Kinali Z. 2017. Phytoplankton dynamics and structure, and ecological status estimation by the Q assemblage index: a comparative analysis in two shallow Mediterranean lakes. *Turkish Journal of Botany* 41:25-36.

Padisák J, Borics G, Fehér G, Grigorszky I, Oldal I, Schmidt A, Zámóné-Doma Z. 2003. Dominant species, functional assemblages and frequency of equilibrium phases in late summer phytoplankton assemblages in Hungarian small shallow lakes. *Hydrobiologia* 502:157-168.

Padisák J, Borics G, Grigorszky I, Soróczki-Pintér É. 2006. Use of phytoplankton assemblages for monitoring ecological status of lakes within the Water Framework Directive: the assemblage index. *Hydrobiologia* 553:1-14.

Padisák J, Crossetti LO, Naselli-Flores L. 2009. Use and misuse in the application of the phytoplankton functional classification: A critical review with updates. *Hydrobiologia* 621:1-19.

Padisák J. 1992. Spatial and temporal scales in phytoplankton ecology. *Abstracta Botanica* 16:15-23.

Padisák J. 1997. *Cylindrospermopsis raciborskii* (Woloszynska) Seenayya et Subba Raju, an expanding, highly adaptive cyanobacterium: worldwide distribution and review of its ecology. *Archiv für Hydrobiologie Supplementband Monographische Beiträge* 107 (4):563-593.

Rangel LM, Silva LH, Arcifa MS, Peticarrari A. 2009. Driving forces of the diel distribution of phytoplankton functional groups in a shallow tropical lake (Lake Monte Alegre, Southeast Brazil). *Brazilian Journal of Biology* 69 (1):75-85.

Reynolds CS, Huszar V, Kruk C, Naselli-Flores L, Melo S. 2002. Towards a functional classification of the freshwater phytoplankton. *Journal of Plankton Research* 24 (5):417-428.

Reynolds CS. 1993. Scales of disturbance and their role in plankton ecology. *Hydrobiologia* 249:157-171.

Reynolds CS. 2006. *The Ecology of Phytoplankton*. Cambridge University Press, 535 pp.

Roll YV. 1961. Fitoplankton sovětského ústí Dunaje, jeho rukavce a záliv. [Phytoplankton of the soviet part of the Danube mouth.] *Tr. In-ta higrobiol. Akad. Nauk USSR* 36 (in Russian)

Rott E. 1981. Some results from phytoplankton counting intercalibration. *Swiss Journal of Hydrology* 43:34-62.

Salmaso N, Padisák J. 2007. Morpho-Functional Groups and phytoplankton development in two deep lakes (Lake Garda, Italy and Lake Stechlin, Germany). *Hydrobiologia* 578 (1):97-112.

Santana LM, Crossetti LO, Ferragut C. 2017. Ecological status assessment of tropical reservoirs through the assemblage index of phytoplankton functional groups. *Brazilian Journal of Botany* 40 (3):695-704.

Schwarz U. 2005 Landschaftsökologische Charakterisierung des Kopački Rit unter besonderer Berücksichtigung von Flusslandschaftsformen sowie deren Genese und Typologie. Dissertation, University of Wien.

Schwarz U. 2010. *Assessment of the restoration potential along the Danube and main tributaries*. Prepared for WWF International Danube Carpathian Programme (DCP), Vienna, 58 pp.

- Silva APC, Costa IAS da. 2015. Biomonitoring ecological status of two reservoirs of the Brazilian semi-arid using phytoplankton assemblages (Q index). *Acta Limnologica Brasiliensia* 27 (1):1-14.
- Sournia A. 1978. *Phytoplankton manual*. UNESCO: Paris.
- Stanković I, Vlahović T, Gligora Udovič M, Várbió G, Borics G. 2012. Phytoplankton functional and morpho-functional approach in large floodplain rivers. *Hydrobiologia* 698:217-231.
- Stević F, Mihaljević M, Horvatić J. 2005. Interactions between microphytoplankton of the Danube, its sidearms and wetlands (1426 - 1388 r. Km, Croatia). *Periodicum Biologorum* 107:299-304.
- Stević F, Mihaljević M, Špoljarić D. 2013. Changes of phytoplankton functional groups in a floodplain lake associated with hydrological perturbations. *Hydrobiologia* 709:143-158.
- Stoyneva MP. 2003. Steady-state phytoplankton assemblages in shallow Bulgarian wetlands. *Hydrobiologia* 502:169-176.
- Strickland JDH, Parson TR. 1972. A practical handbook of seawater analysis. *Fisheries Research Board of Canada Bulletin* 167:1-310.
- Tadić L, Bonacci O, Dadić T. 2014. Dynamics of the Kopački Rit (Croatia) wetland floodplain water regime. *Environmental Earth Sciences* 71 (8):3559-3570.
- Teneva I, Gecheva G, Cheshmedijev S, Stoyanov P, Mladenov R, Belkinova D. 2014. Ecological status assessment of Skalenski Lakes (Bulgaria). *Biotechnology & Biotechnological Equipment* 28 (1):82-95.
- Tolotti M, Boscaini A, Salmaso N. 2010. Comparative analysis of phytoplankton patterns in two modified lakes with contrasting hydrological features. *Aquatic Sciences* 72:213-226.
- UNESCO 1966. Determination of phytoplankton pigments in sea water. Report of SCOR-UNESCO Working Group 17. *Monographs on Oceanographic Methodology* 1, 69.
- Utermöhl H. 1958. Zur Vervollkommnung der quantitative Phytoplankton-Methodik. *Mitteilungen der international Vereinigung für theoretische und angewandte Limnologie* 9:1-38.

- Vidaković J, Bogut I. 2007. Periphyton nematode assemblages in association with *Myriophyllum spicatum* L. in Lake Sakadaš, Croatia. *Russian Journal of Nematology* 15:79-88.
- Vieira PCS, Cardoso MML, Costa IAS da. 2015. Vertical and temporal dynamics of phytoplanktonic associations and the application of index assembly in tropical semi-arid eutrophic reservoir, northeastern Brazil. *Acta Limnologica Brasiliensia* 27 (1):130-144.
- Wang L, Cai Q, Tan L, Kong L. 2011. Phytoplankton development and ecological status during a cyanobacterial bloom in a tributary bay of the Three Gorges Reservoir, China. *Science of the Total Environment* 409:3820-3828.
- WFD, 2000. Directive of the European Parliament and of the Council 2000/60/EC. Establishing a framework for community action in the field of water policy. *Official Journal of the European Union* L327:1-72.
- Xiao LJ, Wang T, Hu R, Han BP, Wang S, Qian X, Padisák J. 2011. Succession of phytoplankton functional groups regulated by monsoonal hydrology in a large canyon-shaped reservoir. *Water Research* 45:5099-5109.
- Yang Y, Stenger-Kovács C, Padisák J, Pettersson K. 2016. Effects of winter severity on spring phytoplankton development in a temperate lake (Lake Erken, Sweden). *Hydrobiologia* 780 (1):47-57.
- Yang Y. 2015. Phytoplankton and physical disturbance. Seasonal dynamics in temperate Lake Erken, Sweden. *Digital Comprehensive Summaries of Uppsala Dissertations from the Faculty of Science and Technology* 1292. Uppsala: Acta Universitatis Upsaliensis. 43pp.

7. APPENDIX

Appendix 1. Biomass of phytoplankton with pertaining functional group in the period July 2011-October 2012.

		Biomass (mg L ⁻¹)																	
		2011						2012											
List of species	FG	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct			
CYANOBACTERIA																			
<i>Anabaena</i> sp.	H1	0.26						<0.005			0.06								
<i>Anabaenopsis arnoldii</i> Aptekar	H1	0.29	0.29										0.38			0.61			
<i>Aphanizomenon flosaquae</i> Ralfs ex Bornet & Flahault	H1	3.79				0.29		0.05			0.12			0.09	3.30	0.69			
<i>Cuspidothrix issatschenkoii</i> (Usachev) P.Rajaniemi, Komárek, R.Willame, P. Hrouzek, K.Kastovská, L.Hoffmann & K.Sivonen	H1	0.25	31.85										0.07			2.56	0.59		
<i>Dolichospermum sigmoideum</i> (Nygaard) Wacklin, L.Hoffmann & Komárek	H1	3.17													4.63				
<i>Dolichospermum solitarium</i> (Klebahn) Wacklin, L.Hoffmann & Komárek	H1	1.37	0.46													0.51	0.30		
<i>Dolichospermum spiroides</i> (Klebhan) Wacklin, L.Hoffmann & Komárek	H1	0.73																	
<i>Aphanocapsa delicatissima</i> West & G.S.West	K					1.73	0.01							0.04					
<i>Aphanocapsa planctonica</i> (G.M.Smith) Komárek & Anagnostidis	K	0.08	0.12	0.20	0.05							0.02	0.07	0.04	0.16	0.02	0.09		
<i>Cyanothece aeruginosa</i> (Nägeli) Komárek	K	0.12																	
<i>Chroococcus distans</i> (G.M.Smith) Komárková-Legnerová & Cronberg	L ₀													0.13					
<i>Chroococcus minutus</i> (Kützing) Nägeli	L ₀	0.03					0.04	0.03	0.01	0.02	0.02	0.06	0.03	0.06	0.07	0.01	0.01		
<i>Chroococcus turgidus</i> (Kützing) Nägeli	L ₀	0.07	0.06			0.03			0.03				0.15						
<i>Gomphosphaeria aponina</i> Kützing	L ₀	0.24	0.38			0.01			0.02				0.03	0.08	0.16				
<i>Gomphosphaeria</i> sp.	L ₀	0.07																	
<i>Limnococcus limneticus</i> (Lemmermann) Komárková, Jezberová, O.Komárek & Zapomelová	L ₀	0.02																	
<i>Merismopedia elegans</i> A.Braun ex Kützing	L ₀	2.19			0.99									0.86			0.57		
<i>Merismopedia glauca</i> (Ehrenberg) Kützing	L ₀	0.07																	
<i>Merismopedia punctata</i> Meyen	L ₀	0.04	0.35	0.43							0.01			0.03					
<i>Merismopedia tenuissima</i> Lemmermann	L ₀													0.06					

List of species		FG	Biomass (mg L ⁻¹)																
			2011						2012										
			Jul	Aug	Sep	Oct	Nov	Dec	Jan	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct		
<i>Snowella lacustris</i> (Chodat) Komárek & Hindák	L ₀	0.06						0.02											
<i>Woronichinia fusca</i> (Skuja) Komárek & Hindák	L ₀	0.18	0.12																
<i>Microcystis aeruginosa</i> (Kützing) Kützing	M	4.32																	
<i>Microcystis viridis</i> (A.Braun) Lemmermann	M	0.15																	
<i>Kamptonema formosum</i> (Bory ex Gomont) Strunecký, Komárek & J.Smarda	S1			2.34	0.26	0.17		0.03	0.79										
<i>Leptolyngbya fragilis</i> (Gomont) Anagnostidis & Komárek	S1			1.62	0.63	0.20	0.37	0.01	0.09	0.08	0.03	0.12	0.09	0.74	0.53	0.75			
<i>Limnothrix redekei</i> (Goor) Meffert	S1						0.49	0.47	0.23	1.05	0.58								
<i>Phormidium</i> sp.	S1	0.56	0.19																
<i>Planktolingbya limnetica</i> (Lemmermann) Komárková-Legnerová & Cronberg	S1		0.29																
<i>Planktothrix agardhii</i> (Gomont) Anagnostidis & Komárek	S1	3.74	7.71	60.19	26.78	1.91	0.32	0.22	0.11	0.36	0.37	5.80					1.12	0.55	
<i>Pseudanabaena limnetica</i> (Lemmermann) Komárek	S1	0.36	1.41	4.31	0.31	0.04	0.81	0.23	0.33	0.50	0.41	0.11	0.01	0.18	1.48	0.91			
<i>Romeria elegans</i> (Woloszynska) Geitler	S1			0.07	0.07			<0.005		0.02	0.03	0.03		1.16	0.65	0.11			
<i>Cylindrospermopsis raciborskii</i> (Woloszynska) Seenayya & Subba Raju	S _N			8.56														0.09	

EUGLENOPHYTA

<i>Euglena texta</i> (Dujardin) Hübner	W1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
--	----	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

		Biomass (mg L ⁻¹)														
		2011						2012								
List of species	FG	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct
<i>Trachelomonas hispida</i> (Perty) F.Stein	W2	0.69														
<i>Trachelomonas oblonga</i> Lemmermann	W2	1.01		0.32	0.69			0.02		0.15					0.30	0.15
<i>Trachelomonas planctonica</i> Svirenko	W2	1.03														
<i>Trachelomonas</i> sp.	W2	0.12														
<i>Trachelomonas volvocina</i> (Ehrenberg) Ehrenberg	W2			0.23	0.92											

PYRROPHYTA

<i>Parvodinium inconspicuum</i> (Lemmermann) S.Carty	L ₀	1.35														
<i>Parvodinium pusillum</i> (Penard) S.Carty	L ₀		0.10													
<i>Peridinium aciculiferum</i> Lemmermann	L ₀	1.16								0.61						
<i>Peridinium bipes</i> Stein	L ₀															0.68
<i>Peridinium cinctum</i> (O.F.Müller) Ehrenberg	L ₀	3.47												1.51	0.69	
<i>Peridinium</i> sp.	L ₀												0.14			
<i>Glenodinium</i> sp.	Y	0.31													0.75	0.37

CRYPTOPHYTA

<i>Rhodomonas lacustris</i> Pascher & Ruttner	X2	0.28	0.15	0.85	1.69	0.70	0.32	0.09	0.39	0.69	0.61	0.40	0.02	1.16	0.59	0.94
<i>Rhodomonas</i> sp.	X2						0.62	0.07	0.41	0.63						
<i>Cryptomonas erosa</i> Ehrenberg	Y			1.64	6.05	0.30	0.17	0.08	0.75	0.70	1.21	1.13	0.41	3.00	0.79	0.48
<i>Cryptomonas ovata</i> Ehrenberg	Y			14.54		2.23	1.27	0.72	2.23	1.87	0.64	2.43	0.69	5.99	3.49	1.62
<i>Cryptomonas</i> sp.	Y	14.27	7.23													

CHRYSTOPHYTA

Xanthophyceae																
<i>Centritractus belonophorus</i> (Schmidle) Lemmermann	J												0.03			0.02
<i>Goniochloris mutica</i> (A.Braun) Fott	J				0.03								0.01			

		Biomass (mg L ⁻¹)																		
		2011						2012												
List of species	FG	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct				
<i>Ophiocytium capitatum</i> var. <i>longispinum</i> (Möbius) Lemmermann	J	0.17																		
<i>Ophiocytium capitatum</i> Wolle	J							0.02			0.04	0.09								
<i>Tetraplektron tribulus</i> (Pascher) Lobelich	J																0.02			
Chrysophyceae																				
<i>Dinobryon divergens</i> O.E.Imhof	E	0.18						0.15		0.01		0.13								
<i>Salpingoeca frequentissima</i> (Zacharias) Lemmermann	E																0.01			
<i>Synura uvella</i> Ehrenberg	W _s							0.54												
<i>Bitrichia danubiensis</i> Juri	X2																0.03			
<i>Kephyrion rubi-claustri</i> Conrad	X2							0.05	0.02	0.22	0.01					0.01				
<i>Kephyrion</i> sp.	X2							0.01	<0.005											
<i>Chrysococcus rufescens</i> Klebs	X3			0.41	1.10	0.02	0.16	0.03	0.29	0.35	0.13	0.08	0.05	0.27	0.15	0.15				
Bacillariophyceae																				
<i>Acanthoceras zachariasii</i> (Brun) Simonsen	A	0.52																		
<i>Cyclotella</i> sp.	A							0.21						0.21						
<i>Lindavia comta</i> (Kützing) Nakov, Gullory, Julius, Theriot & Alverson	B							0.04												
<i>Asterionella formosa</i> Hassall	C	0.16	0.16					0.02	0.19		0.25	0.08								
<i>Cyclotella meneghiniana</i> Kützing	C	0.55	0.13			2.48	23.17	0.30	0.22	0.36	0.44		1.17	0.34	0.79	0.73 0.74				
<i>Fragilaria acus</i> (Kützing) Lange-Bertalot in Krammer & Lange-Bertalot	D	0.10	0.02	0.66	0.08	1.53	0.09	0.19	0.08	0.36	0.09			0.10	1.85	0.06				
<i>Nitzschia acicularis</i> (Kützing) W.Smith	D			0.12	0.02	0.02		0.01	0.01		0.02			0.03						
<i>Nitzschia fusiformis</i> Grunow	D	0.08																		
<i>Nitzschia holsatica</i> Hustedt	D																0.13	0.18	0.10	0.01
<i>Nitzschia palea</i> (Kützing) W.Smith	D	0.03						0.01								0.16	0.32			
<i>Nitzschia</i> sp. 1	D	0.03	0.01																	
<i>Nitzschia</i> sp. 2	D	0.09	0.05																	

		Biomass (mg L ⁻¹)															
		2011						2012									
List of species	FG	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	
<i>Skeletonema potamos</i> (C.I.Weber) Hasle in Hasle & Evensen	D	0.01						0.02									
<i>Stephanodiscus hantzschii</i> Grunow	D	2.88	4.74	1.27	6.01	7.70	2.09	0.11	0.34			0.91	0.14		0.24	1.16	
<i>Ulnaria ulna</i> (Nitzsch) Compère	D	2.34	16.41	1.95		1.42			1.18	4.78		2.43	1.53	7.13	0.47		
<i>Amphora ovalis</i> (Kützing) Kützing	MP	3.64	0.23														
<i>Cocconeis placentula</i> Ehrenberg	MP	0.35															
<i>Nitzschia sigmoidea</i> (Nitzsch) W.Smith	MP	0.51															
<i>Aulacoseira granulata</i> (Ehrenberg) Simonsen	P	1.51	1.08		0.52	1.04						1.05					
<i>Aulacoseira granulata</i> var. <i>angustissima</i> (Otto Müller) Simonsen	P					0.88	0.20		0.07		0.11	0.52	0.42	0.51			
<i>Fragilaria capucina</i> Desmazières	P	0.26															
<i>Cymbella</i> sp.	T _B	0.11															
<i>Cymbella tumida</i> (Brébisson) Van Heurck	T _B	0.11															
<i>Gomphonema parvulus</i> (Lange-Bertalot & E.Reichardt) Lange-Bertalot & E.Reichardt	T _B	0.05															
<i>Melosira varians</i> C.Agardh	T _B	0.38															
<i>Navicula</i> sp. 1	T _B	0.03	0.03					0.01					0.05				
<i>Navicula</i> sp. 2	T _B	0.09	0.42														
<i>Navicula</i> sp. 3	T _B	0.05															
CHLOROPHYTA																	
<i>Chlorobotrys</i> sp.	F	0.22	0.13										0.04	0.08	0.14		
<i>Dictyosphaerium ehrenbergianum</i> Nägeli	F	0.12	0.03	0.03										0.02			
<i>Hindakia tetrachotoma</i> (Printz) C.Bock, Pröschold & Krienitz	F	0.15	0.03					0.01									
<i>Keratococcus bicaudatus</i> (A.Braun ex Rabenhorst) J.B.Petersen	F							0.11									
<i>Kirchneriella lunaris</i> (Kirchner) Möbius	F															<0.005	
<i>Kirchneriella</i> sp.	F	0.03															
<i>Micractinium bornhemiense</i> (W.Conrad) Korshikov	F	2.08						0.45									

		Biomass (mg L ⁻¹)														
		2011						2012								
List of species	FG	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct
<i>Micractinium pusillum</i> Fresenius	F	0.29	0.38			0.35	0.26					0.13	0.13			0.45
<i>Mucidosphaerium pulchellum</i> (H.C.Wood) C.Bock, Proschold & Krienitz	F	0.05	0.02									0.05	0.04			
<i>Oocystis lacustris</i> Chodat	F	0.31	0.08													
<i>Oocystis</i> sp.	F	0.12	0.12													
<i>Treubaria planctonica</i> (G.M.Smith) Korshikov	F													0.03		
<i>Treubaria schmidlei</i> (Schröder) Fott & Kovácik	F														0.02	0.02
<i>Actinastrum hantzschii</i> Lagerheim	J	0.17	0.11	0.29	0.16							0.08	0.37	0.30	0.03	0.07
<i>Chlorotetraedron incus</i> (Teiling) Komárek & Kovácik	J		0.03	0.07							0.01	0.02	0.02			0.02
<i>Coelastrum astroideum</i> De Notaris	J	1.10	0.18	0.46		0.17					0.10					0.11
<i>Coelastrum microporum</i> Nägeli	J	0.43	0.65						0.07		0.46	0.15	0.18		1.04	
<i>Crucigenia quadrata</i> Morren	J											0.01	0.01		0.14	0.04
<i>Crucigenia tetrapedia</i> (Kirchner) Kuntze	J	1.01	0.13	0.07	0.04	0.01	0.01		<0.005		0.02					
<i>Desmodesmus arthrodesmiformis</i> (Schröder) S.S.An, Friedl & E.Hegewald	J				0.09							0.02	0.07	0.04		
<i>Desmodesmus bicaudatus</i> (Dedusenko) P.M.Tsarenko	J	0.07			0.05	0.03					0.02	0.06	0.04		0.20	0.02
<i>Desmodesmus denticulatus</i> (Lagerheim) S.S.An, T.Friedl & E.Hegewald	J				0.18											0.02
<i>Desmodesmus intermedius</i> (Chodat) E.Hegewald	J		0.16													
<i>Desmodesmus opoliensis</i> (P.G.Richter) E.Hegewald	J	0.49										0.06				0.05
<i>Desmodesmus spinosus</i> (Chodat) E.Hegewald	J							0.01		0.02	0.08		0.03			
<i>Franceia ovalis</i> (Francé) Lemmermann	J											0.03				
<i>Golenkinia radiata</i> Chodat	J		0.16		0.21	0.28	0.05					0.03			0.14	0.37
<i>Lagerheimia ciliata</i> (Lagerheim) Chodat	J		0.06													
<i>Lagerheimia genevensis</i> (Chodat) Chodat	J				0.01		<0.005	<0.005	<0.005	<0.005	<0.005	<0.005				
<i>Lagerheimia longiseta</i> (Lemmermann) Printz	J		0.01													<0.005
<i>Lagerheimia wratislaviensis</i> f. <i>heterospina</i> Hortobágyi	J		0.01													
<i>Monactinus simplex</i> (Meyen) Corda	J				0.14								0.03		0.06	

		Biomass (mg L ⁻¹)																	
		2011						2012											
List of species	FG	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct			
<i>Pediastrum boryanum</i> var. <i>cornutum</i> (Raciborski) Sulek	J							0.05											
<i>Pediastrum duplex</i> Meyen	J	0.20											0.06	0.12					
<i>Pseudopediastrum boryanum</i> (Turpin) E.Hegewald	J	0.33			0.18			0.05					0.08			0.08			
<i>Pseudotetraëdron neglectum</i> Pascher	J															0.02			
<i>Scenedesmus ecornis</i> (Ehrenberg) Chodat	J						0.03	0.02											
<i>Scenedesmus obtusus</i> f. <i>disciformis</i> (Chodat) Compère	J															0.02			
<i>Scenedesmus obtusus</i> Meyen	J												0.02	0.03					
<i>Scenedesmus quadricauda</i> (Turpin) Brébisson	J	1.01	0.40	0.76	0.55	0.64		0.06	0.03	0.06	0.43	0.17	0.31	0.26	0.18	0.12			
<i>Scenedesmus</i> sp.	J	0.06																	
<i>Stauridium tetras</i> (Ehrenberg) E.Hegewald	J												0.03	0.03					
<i>Tetradesmus lagerheimii</i> M.J.Wynne & Guiry	J	0.42	0.14	0.19		0.25		1.02				0.36	0.16	0.08					
<i>Tetradesmus obliquus</i> (Turpin) M.J.Wynne	J	0.14																	
<i>Tetraëdron minimum</i> (A.Braun) Hansgirg	J	0.03	0.02			0.01	<0.005	<0.005	0.01			0.01				0.02			
<i>Tetraëdron trigonum</i> (Nägeli) Hansgirg	J						0.02												
<i>Tetrastrum glabrum</i> (Y.V.Roll) Ahlstrom & Tiffany	J	0.03			0.01			<0.005	0.01	0.01			0.02	0.01		0.02			
<i>Tetrastrum staurogeniiforme</i> (Schröder) Lemmermann	J	0.02							<0.005		0.01	0.01	0.01	0.04	0.01				
<i>Treubaria triappendiculata</i> C.Bernard	J	0.04																	
<i>Willea apiculata</i> (Lemmermann) D.M.John, M.J.Wynne & P.M.Tsarenko	J	0.09																	
<i>Willea rectangularis</i> (A.Braun) D.M.John, M.J.Wynne & P.M.Tsarenko	J							0.01						0.06	0.13		0.05		
<i>Cosmarium laeve</i> Rabenhorst	N															0.08			
<i>Cosmarium phaseolus</i> Brébisson ex Ralfs	N	0.32	0.32				0.22									0.57			
<i>Closteriopsis acicularis</i> (Chodat) J.H.Belcher & Swale	P	0.09			0.05	0.03	0.03	0.01				<0.005	0.08		0.10	0.01			
<i>Closterium diana</i> e Ehrenberg ex Ralfs	P	0.28																	
<i>Closterium ehrenbergii</i> Meneghini ex Ralfs	P	0.28																	
<i>Closterium gracile</i> Brébisson ex Ralfs	P	0.28	0.28															0.37	
<i>Closterium intermedium</i> Ralfs	P															0.33			

		Biomass (mg L ⁻¹)																		
		2011						2012												
List of species	FG	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct				
<i>Closterium kuetzingii</i> Brébisson	P	0.34																		
<i>Closterium limneticum</i> Lemmermann	P	0.40						0.02												
<i>Staurastrum</i> sp.	P															0.28				
<i>Staurastrum tetracerum</i> Ralfs ex Ralfs	P															0.28				
<i>Binuclearia lauterbornii</i> (Schmidle) Proschkina-Lavrenko	T	0.19														0.69	2.29	0.21		
<i>Mougeotia</i> sp.	T	0.35		4.83		0.48										0.10	0.69	0.11	0.21	
<i>Ankistrodesmus arcuatus</i> Korshikov	X1		0.01	0.02	0.02	0.01	0.02	0.03	0.04	0.07		0.01		0.01	<0.005					
<i>Ankistrodesmus falcatus</i> (Corda) Ralfs	X1	0.01																		
<i>Ankistrodesmus fusiformis</i> Corda	X1	0.04					0.01	0.01								0.03	0.01			
<i>Chlorolobion braunii</i> (Nägeli) Komárek	X1															<0.005	0.01	0.03		
<i>Messastrum gracile</i> (Reinsch) T.S.Garcia	X1	0.03				0.01										0.01				
<i>Monoraphidium contortum</i> (Thuret) Komárková-Legnerová	X1	0.09	0.22	0.01	0.13	0.02	0.05	0.01	0.02	0.14	0.24	0.08	0.07	0.08	0.02	0.03				
<i>Monoraphidium convolutum</i> (Corda) Komárková-Legnerová	X1															0.02				
<i>Monoraphidium griffithii</i> (Berkeley) Komárková-Legnerová	X1	0.07															0.02	0.02		
<i>Monoraphidium irregulare</i> (G.M.Smith) Komárková-Legnerová	X1	0.02	1.30	0.74							0.05	0.06							0.10	0.12
<i>Monoraphidium komarkovae</i> Nygaard	X1	0.02	0.01																	
<i>Monoraphidium minutum</i> (Nägeli) Komárková-Legnerová	X1	0.02	0.01															0.02		
<i>Polyedriopsis spinulosa</i> (Schmidle) Schmidle	X1	0.14						0.17												
<i>Pseudodidymocystis inconspicua</i> (Korshikov) Hindák	X1	0.06	0.02																	
<i>Pseudodidymocystis planctonica</i> (Korshikov) E.Hegewald & Deason	X1	0.02						<0.005	0.11	0.01	0.05	0.07	0.16	0.14	0.13	0.05				
<i>Pseudosphaerocystis lacustris</i> (Lemmermann) Nováková	X1	0.04																		
<i>Schroederia spiralis</i> (Printz) Korshikov	X1	0.76						0.03									0.10	0.58	0.05	
<i>Selenastrum bibraianum</i> Reinsch	X1	0.10														0.01				
<i>Carteria pseudoglobosa</i> Ettl	X2							0.16								0.04				
<i>Carteria</i> sp.	X2	0.48	0.79							0.12	0.25						0.48		1.50	

		Biomass (mg L ⁻¹)														
		2011						2012								
List of species	FG	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct
<i>Chlamydomonas incerta</i> Pascher	X2	0.12														
<i>Chlamydomonas pertusa</i> Chodat	X2							0.09			0.10	0.04		0.47		
<i>Chlamydomonas</i> sp.	X2	3.13	3.18													
<i>Coccomonas orbicularis</i> Stein	X2	0.04														
<i>Platymonas cordiformis</i> Korshikov	X2							0.23								
<i>Schroederia setigera</i> (Schröder) Lemmermann	X3										0.21	0.10	0.33			
<i>Koliella longiseta</i> (Vischer) Hindák	X3	0.02		0.03		0.13	0.04	0.05	0.08	0.15	0.15	<0.005		0.66	0.02	0.02
Total biomass (mg L ⁻¹)		65.07	52.13	146.21	55.65	47.59	8.16	3.13	7.99	13.44	9.02	13.32	8.63	48.53	19.94	17.58
Number of species		85	63	39	44	39	33	37	31	26	40	52	59	44	58	61